

Extending Growth Mixture Models Using Continuous Non-Elliptical Distributions

Yuhong Wei* Emilie Shireman†
 Paul D. McNicholas* Douglas L. Steinley†

Abstract

Growth mixture models (GMMs) incorporate both conventional random effects growth modeling and latent trajectory classes as in finite mixture modeling; therefore, they offer a way to handle the unobserved heterogeneity between subjects in their development. GMMs with Gaussian random effects dominate the literature. When the data are asymmetric and/or have heavier tails, more than one latent class is required to capture the observed variable distribution. Therefore, a GMM with continuous non-elliptical distributions is proposed to capture skewness and heavier tails in the data set. Specifically, multivariate skew- t distributions and generalized hyperbolic distributions are introduced to extend GMMs. When extending GMMs, four statistical models are considered with differing distributions of measurement errors and random effects. The mathematical development of a GMM with non-elliptical distributions relies on its relationship with the generalized inverse Gaussian distribution. Parameter estimation is outlined within the expectation-maximization framework before the performance of our GMM with non-elliptical distributions is illustrated on simulated and real data.

1 Introduction

Many longitudinal studies focus on investigating how individuals change over time with respect to a characteristic that is measured repeatedly for each participant. Conventional random effect growth modeling has provided a number of tools for modeling intra-individual change and inter-individual differences in change (e.g., Laird and Ware, 1982; Bryk and Raudenbush, 1987, 1992; McArdle and Epstein, 1987; Singer and Willett, 2003), where within-class changes are described as a function of time and between-class changes are described by random effects and coefficients. Conventional random effects models provide a basis for formulating growth mixture models (GMMs) for longitudinal data. In the latent variable

*Department of Mathematics & Statistics, McMaster University, Hamilton, ON, Canada.

†Department of Psychological Sciences, University of Missouri, Columbia, MO, U.S.

framework, a quadratic random effects growth model with a continuous outcome y_{it} for individual i at time t , and time-invariant covariates x , is specified according to

$$y_{it} = \eta_{0i} + \eta_{1i}(a_t - a_0) + \eta_{2i}(a_t - a_0)^2 + \epsilon_{it}, \quad (1)$$

$$\begin{aligned} \eta_{0i} &= \alpha_0 + \gamma_0 x_i + \zeta_{0i}, \\ \eta_{1i} &= \alpha_1 + \gamma_1 x_i + \zeta_{1i}, \\ \eta_{2i} &= \alpha_2 + \gamma_2 x_i + \zeta_{2i}, \end{aligned} \quad (2)$$

where a_t are time scores ($t = 1, 2, \dots, T$) centred at a_0 , η_{0i} is the random intercept, η_{1i} is the random linear slope, η_{2i} is the quadratic growth rate, and ϵ_i and ζ_i are normally distributed residuals, i.e., $\epsilon_i \sim \mathcal{N}(\mathbf{0}, \mathbf{\Theta})$ and $\zeta_i \sim \mathcal{N}(\mathbf{0}, \mathbf{\Psi})$. Formally, η_{0i} , η_{1i} , and η_{2i} are continuous latent variables (called the growth factors) representing the growth patterns, the α_k are the mean parameters for the growth factors if there are no covariates $\mathbf{x}_i$, and the γ_k are the effects of covariates $\mathbf{x}_i$ on the growth factors. In conventional growth modeling applications, the individual growth parameters (e.g., individual intercept and slope factors) are usually assumed to be identically distributed, i.e., drawn from a single homogeneous population. However, we are often interested in and deal with samples from multiple populations and, in most cases, the class memberships are either unknown or unobserved.

Simultaneous modeling of change over time and unobserved multiple populations (heterogeneity) in the data can be accommodated using GMMs and latent class growth analysis (LCGA), wherein parameters describing growth patterns are estimated and each individual's most likely class membership is obtained via maximum *a posteriori* (MAP) probabilities. GMMs were introduced by Verbeke and Lesaffre (1996) and then extended by Muthén and Shedden (1999), Muthén (2004), and Muthén and Asparouhov (2008). For convenience, define c_{ik} so that $c_{ik} = 1$ if individual i falls in class k and $c_{ik} = 0$ otherwise. The quadratic growth model of (1) and (2) can be extended to a simple GMM in class k ($k = 1, 2, \dots, K$) via

$$Y_{it} \mid c_{ik} = 1 = \eta_{0i} + \eta_{1i}(a_t - a_0) + \eta_{2i}(a_t - a_0)^2 + \epsilon_{it}, \quad (3)$$

$$\begin{aligned} \eta_{0i} \mid c_{ik} = 1 &= \alpha_{0k} + \gamma_0 x_i + \zeta_{0i}, \\ \eta_{1i} \mid c_{ik} = 1 &= \alpha_{1k} + \gamma_1 x_i + \zeta_{1i}, \\ \eta_{2i} \mid c_{ik} = 1 &= \alpha_{2k} + \gamma_2 x_i + \zeta_{2i}, \end{aligned} \quad (4)$$

where $\boldsymbol{\alpha}_k = (\alpha_{0k}, \alpha_{1k}, \alpha_{2k})$ parameters vary across classes to capture different trajectories; the parameters $\gamma_0, \gamma_1, \gamma_2$, remain the same across classes but this could be relaxed to allow variation across classes with respect to how a covariate affects the growth factors; and ϵ_i and ζ_i are still normally distributed residuals but with class-specific covariance matrices, i.e., $\epsilon_i \sim \mathcal{N}(\mathbf{0}, \mathbf{\Theta}_k)$ and $\zeta_i \sim \mathcal{N}(\mathbf{0}, \mathbf{\Psi}_k)$. The latent class growth analysis (LCGA) developed by Nagin and colleagues (Nagin and Land, 1993; Nagin, 1999, 2005) can be thought of as one special case of GMMs in the sense that it assumes zero within-class growth factor variances, i.e., $\mathbf{\Psi}_k = \mathbf{0}$ for $k = 1, 2, \dots, K$.

One common fundamental assumption for GMMs is that model errors are normally distributed. However, when the data are asymmetric and/or have heavier tails (skewed), more than one latent class is required to capture the observed variable distribution; thus, interpreting mixture components as subpopulations will lead to overestimation of the subpopulations. Being able to relax the normality assumption for asymmetry and skewness, Muthén and Asparouhov (2015) developed a GMM with a normally distributed model error and “classical” multivariate skew-t (cMST) random effects, i.e., $\epsilon_i \sim \mathcal{N}(\mathbf{0}, \Theta_k)$ and $\zeta_i \sim \text{MST}(\mathbf{0}, \Psi_k)$. An alternative specification of GMMs (called nonlinear mixed effect mixture models) was developed by Lu and Huang (2014) within a Bayesian framework, wherein ϵ_i is assumed to follow a cMST while ζ_i remains normally distributed. Note that the “classical” formulation of the multivariate skew-t distribution is that given by Azzalini and Valle (1996) and Azzalini and Capitanio (1999). In this paper, we outline a more general extension of GMMs to the generalized hyperbolic distribution while also considering the formulation of the multivariate skew-t distribution that arises as its special and limiting case (Section 2). The advantage of the generalized hyperbolic distribution is its flexibility. Many other well-known distributions are special or limiting cases of the generalized hyperbolic distribution; please refer to McNeil et al. (2005) for details on a variety of limiting cases of the generalized hyperbolic distribution.

The remainder of this article is laid out as follows. In Section 2, we go through some important background material on the generalized hyperbolic distribution as well as a special and limiting case that gives a formulation of the multivariate skew-t distribution. In Section 3, we outline the extension of GMMs to generalized hyperbolic and multivariate skew-t distributions, respectively. Section 4 presents an expectation-maximization (EM) algorithm for obtaining the maximum likelihood (ML) estimates of model parameters. Then, our approach is illustrated on simulated and real data (Section 5). The paper concludes with some discussion and suggestions for future work (Section 6).

2 Background

2.1 Generalized hyperbolic distribution

A p -dimensional multivariate generalized hyperbolic distribution, as represented by Browne and McNicholas (2015), has density

$$f(\mathbf{y} \mid \boldsymbol{\vartheta}) = \left[\frac{\omega + \delta(\mathbf{y}, \boldsymbol{\mu} \mid \boldsymbol{\Sigma})}{\omega + \boldsymbol{\beta}' \boldsymbol{\Sigma}^{-1} \boldsymbol{\beta}} \right]^{(\lambda - p/2)/2} \frac{K_{\lambda - p/2}(\sqrt{[\omega + \boldsymbol{\beta}' \boldsymbol{\Sigma}^{-1} \boldsymbol{\beta}] [\omega + \delta(\mathbf{y}, \boldsymbol{\mu} \mid \boldsymbol{\Sigma})]})}{(2\pi)^{p/2} \mid \boldsymbol{\Sigma} \mid^{1/2} K_{\lambda}(\omega) \exp\{-(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} \boldsymbol{\beta}\}}, \quad (5)$$

with index parameter λ , concentration parameter ω , skewness parameter $\boldsymbol{\beta}$, mean vector $\boldsymbol{\mu}$, and scale matrix $\boldsymbol{\Sigma}$. Here, $\delta(\mathbf{y}, \boldsymbol{\mu} \mid \boldsymbol{\Sigma})$ is the squared Mahalanobis distance between $\mathbf{y}$ and $\boldsymbol{\mu}$, i.e., $\delta(\mathbf{y}, \boldsymbol{\mu} \mid \boldsymbol{\Sigma}) = (\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \boldsymbol{\mu})$, $K_{\lambda - p/2}$ and K_{λ} are modified Bessel functions of the third kind with indices $\lambda - p/2$ and λ , respectively, and $\boldsymbol{\vartheta} = (\lambda, \omega, \boldsymbol{\mu}, \boldsymbol{\Sigma}, \boldsymbol{\beta})$ denotes the model

parameters. Herein, let $\mathbf{Y} \sim \text{GHD}_p(\lambda, \omega, \boldsymbol{\mu}, \boldsymbol{\Sigma}, \boldsymbol{\beta})$ represent a p -dimensional generalized hyperbolic random variable $\mathbf{Y}$ with density as per (5). Note that the parameterization in (5) is one of several available for multivariate generalized hyperbolic distributions (cf. McNeil et al., 2005).

By introducing a generalized inverse Gaussian (GIG) random variable $W \sim \mathcal{I}(\omega, \eta, \lambda)$ and a latent multivariate Gaussian random vector $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma})$, then a generalized hyperbolic random variable $\mathbf{Y}$ in (5) can be generated using the relationship

$$\mathbf{Y} = \boldsymbol{\mu} + W\boldsymbol{\beta} + \sqrt{W}\mathbf{Z}, \quad (6)$$

where we fix $\eta = 1$. Herein, the density of $W \sim \mathcal{I}(\omega, \eta, \lambda)$ is

$$h(w \mid \omega, \eta, \lambda) = \frac{(w/\eta)^{\lambda-1}}{2\eta K_\lambda(\omega)} \exp \left\{ -\frac{\omega}{2} \left(\frac{w}{\eta} + \frac{\eta}{w} \right) \right\}. \quad (7)$$

Note that the density in (7) is sometimes a more meaningful reparameterization of the GIG distribution, denoted by $\text{GIG}(\psi, \chi, \lambda)$ (Good, 1953; Barndorff-Nielsen and Halgreen, 1977; Blæsild, 1978; Halgreen, 1979; Jørgensen, 1982). The links between these two parameterizations of the GIG distribution are $\omega = \sqrt{\psi\chi}$ and $\eta = \sqrt{\chi/\psi}$. Extensive details on the derivative of these parameterizations and the relationship in (6) are given by Browne and McNicholas (2015). The GIG distribution has some attractive features, including tractable expected values. Consider $W \sim \mathcal{I}(\omega, \eta, \lambda)$, then the following expected values hold:

$$\begin{aligned} \mathbb{E}[W] &= \eta \frac{K_{\lambda+1}(\omega)}{K_\lambda(\omega)}, \\ \mathbb{E}[1/W] &= \frac{1}{\eta} \frac{K_{\lambda+1}(\omega)}{K_\lambda(\omega)} - \frac{2\lambda}{\omega\eta}, \\ \mathbb{E}[\log W] &= \log \eta + \frac{1}{K_\lambda(\omega)} \frac{\partial}{\partial \lambda} K_\lambda(\omega). \end{aligned} \quad (8)$$

From (6), it follows that $\mathbf{Y} \mid w \sim \mathcal{N}(\boldsymbol{\mu} + w\boldsymbol{\beta}, w\boldsymbol{\Sigma})$. Now, recalling that $W \sim \mathcal{I}(\omega, 1, \lambda)$ and that the unconditional distribution of $\mathbf{Y}$ is generalized hyperbolic, Bayes' theorem gives $W \mid \mathbf{y} \sim \text{GIG}(\omega + \boldsymbol{\beta}'\boldsymbol{\Sigma}^{-1}\boldsymbol{\beta}, \omega + \delta(\mathbf{y}, \boldsymbol{\mu} \mid \boldsymbol{\Sigma}), \lambda - p/2)$. This elegant result will be used to extend GMMs to the generalized hyperbolic distribution.

2.2 Multivariate skew-t distribution

Several alternative formulations of the multivariate skew-t distribution have appeared in the literature (e.g., Sahu et al., 2003; Lin et al., 2014; Lee and McLachlan, 2014; McNeil et al., 2005). Muthén and Asparouhov (2015) developed a GMM with Sahu's version of the restricted multivariate skew-t distribution. The formulation of the multivariate skew-t distribution used herein arises as a special and limiting case of the generalized hyperbolic

distribution by setting $\lambda = -\nu/2$ and $\chi = \nu$, and letting $\psi \rightarrow 0$. This formulation of the multivariate skew-t distribution has been used by Murray et al. (2014) to develop a mixture of skew-t factor analyzer models.

A p -dimensional skew-t random variable $\mathbf{Y}$ has the density function

$$f(\mathbf{y} \mid \boldsymbol{\vartheta}) = \left[\frac{\nu + \delta(\mathbf{y}, \boldsymbol{\mu} \mid \boldsymbol{\Sigma})}{\boldsymbol{\beta}' \boldsymbol{\Sigma}^{-1} \boldsymbol{\beta}} \right]^{(-\nu-p)/4} \frac{\nu^{\nu/2} K_{(-\nu-p)/2}(\sqrt{[\boldsymbol{\beta}' \boldsymbol{\Sigma}^{-1} \boldsymbol{\beta}] [\nu + \delta(\mathbf{y}, \boldsymbol{\mu} \mid \boldsymbol{\Sigma})]})}{(2\pi)^{p/2} |\boldsymbol{\Sigma}|^{1/2} \Gamma(\nu/2) \nu^{1/2} \exp\{-(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} \boldsymbol{\beta}\}}, \quad (9)$$

where $\boldsymbol{\mu}$ is the location parameter, $\boldsymbol{\Sigma}$ is the scale parameter, $\boldsymbol{\beta}$ is the skew parameter, ν is the degree of freedom parameter, and $K_{(-\nu-p)/2}$ and $\delta(\mathbf{y}, \boldsymbol{\mu} \mid \boldsymbol{\Sigma})$ are as defined in (5). We write $\mathbf{Y} \sim \text{GST}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \boldsymbol{\beta}, \nu)$ to denote that the random variable $\mathbf{Y}$ follows the skew-t distribution such that it has the density in (9). Now, $\mathbf{Y} \sim \text{GST}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \boldsymbol{\beta}, \nu)$ can be obtained through the relationship in (6) with $W \sim \text{IG}(\nu/2, \nu/2)$, where $\text{IG}(\cdot)$ denotes the inverse-gamma distribution. We have $\mathbf{Y} \mid w \sim \mathcal{N}(\boldsymbol{\mu} + w\boldsymbol{\beta}, w\boldsymbol{\Sigma})$, and so, from Bayes's theorem, $W \mid \mathbf{y} \sim \text{GIG}(\boldsymbol{\beta}' \boldsymbol{\Sigma}^{-1} \boldsymbol{\beta}, \nu + \delta(\mathbf{y}, \boldsymbol{\mu} \mid \boldsymbol{\Sigma}), -(\nu + p)/2)$.

2.3 The EM Algorithm and its convergence criterion

The EM algorithm (Dempster et al., 1977) is a two-step iterative algorithm for finding maximum likelihood estimates when data are incomplete or treated as such, and is widely used to estimate model parameters in the context of model-based clustering. The E-step computes the expected value of the complete-data log-likelihood given the current model parameters, and the M-step maximizes this expected value with respect to the model parameters. After each E- and M-step, the log-likelihood is driven uphill, and the method iterates towards a maximum until some convergence criterion is satisfied. Many variants of the EM algorithm have been proposed over the years, such as the expectation-conditional-maximization (ECM) algorithm (Meng and Rubin, 1993), the alternating ECM (AECM) algorithm (Meng and Van Dyk, 1997), and Fisher's EM algorithm (Bouveyron and Brunet, 2012). Herein, we will make use of the EM algorithm for parameter estimation.

A stopping criterion based on the Aitken acceleration (Aitken, 1926) is used to determine the convergence of the EM algorithm. The Aitken acceleration at iteration s is

$$a^{(s)} = \frac{l^{(s+1)} - l^{(s)}}{l^{(s)} - l^{(s-1)}}, \quad (10)$$

where $l^{(s)}$ is the log-likelihood value at iteration s . This yields an asymptotic estimate of the log-likelihood at iteration $s + 1$, given by

$$l_{\infty}^{(s+1)} = l^{(s)} + \frac{1}{1 - a^{(s)}}(l^{(s+1)} - l^{(s)})$$

(Böhning et al., 1994; Lindsay, 1995), and the EM algorithm is stopped when $l_{\infty}^{(s+1)} - l^{(s)} < \epsilon$, provided this difference is positive (McNicholas et al., 2010). Note that this criterion is at least as strict as the lack of progress criterion.

2.4 Model selection

In model-based clustering, a penalized log-likelihood-based criterion is typically used to determine the ‘best’ fitting model among a family of models. The most popular such criterion is the Bayesian information criterion (BIC; Schwarz, 1978), which is motivated by the use of Bayes factors (Kass and Raftery, 1995; Kass and Wasserman, 1995). The BIC is defined as

$$\text{BIC} = -2l(\hat{\boldsymbol{\vartheta}}) + \rho \log n, \quad (11)$$

where $\hat{\boldsymbol{\vartheta}}$ is the maximum likelihood estimate of model parameters $\boldsymbol{\vartheta}$, $l(\hat{\boldsymbol{\vartheta}})$ is the maximized log-likelihood, ρ is the number of free parameters, and n is the number of observations. Some theoretical support for use of the BIC in mixture model selection is given by Leroux (1992) and Keribin (2000).

3 Methodology

3.1 Conventional GMM with Gaussian random effects

Suppose a longitudinal study features n subjects and T time points or measurement occasions. For subject i ($i = 1, \dots, n$), let $\mathbf{y}_i$ be a $T \times 1$ vector $\mathbf{y}_i = (y_{i1}, y_{i2}, \dots, y_{iT})'$ where y_{it} represents the outcome on occasion t ($t = 1, \dots, T$), let $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{im})'$ be an $m \times 1$ vector of observed time-invariant covariates, let $\boldsymbol{\eta}_i$ be a $q \times 1$ vector containing q continuous latent variables, and note that $\mathbf{C}_i = (C_{i1}, \dots, C_{iK})'$ has a multinomial distribution, where $C_{ik} = 1$ if individual i is in class k and $C_{ik} = 0$ otherwise. The conventional GMM with Gaussian random effects can be represented using a hierarchical three-level formulation as follows.

At level 1 of the GMM, the continuous outcome variables $\mathbf{Y}_1, \dots, \mathbf{Y}_n$ are related to the continuous latent variables $\boldsymbol{\eta}_1, \dots, \boldsymbol{\eta}_n$ via

$$\mathbf{Y}_i = \boldsymbol{\Lambda}_y \boldsymbol{\eta}_i + \boldsymbol{\epsilon}_i, \quad (12)$$

for $i = 1, \dots, n$, $\boldsymbol{\epsilon}_i$ is a $T \times 1$ vector of residuals or measurement errors that is assumed to follow a multivariate normal distribution $\boldsymbol{\epsilon}_i \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Theta}_k)$, and $\boldsymbol{\Lambda}_y$ is a $T \times q$ design matrix consisting of factor loadings with each column corresponding to specific aspects of change. The matrix $\boldsymbol{\Lambda}_y$ and the vector $\boldsymbol{\eta}_i$ determine the growth trajectory of the model. For instance, when $q = 3$, $\boldsymbol{\eta}_i = (\eta_{0i}, \eta_{1i}, \eta_{2i})$, and $\boldsymbol{\Lambda}_y$ is a $T \times 3$ matrix. Assuming a_t are age-related time scores ($t = 1, 2, \dots, T$) centred at age a_0 , then $\boldsymbol{\Lambda}_y$ is given by

$$\boldsymbol{\Lambda}_y = \begin{pmatrix} 1 & a_1 - a_0 & (a_1 - a_0)^2 \\ 1 & a_2 - a_0 & (a_2 - a_0)^2 \\ \vdots & \vdots & \vdots \\ 1 & a_{T-1} - a_0 & (a_{T-1} - a_0)^2 \end{pmatrix}.$$

At level 2 of the GMM, the continuous latent variables $\boldsymbol{\eta}$ are related to the latent categorical variables $\mathbf{c}$ and to the observed time-invariant covariate vector $\mathbf{x}$ by the relation

$$\boldsymbol{\eta}_i = \boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + \boldsymbol{\zeta}_i, \quad (13)$$

where $\boldsymbol{\alpha}_k$ ($k = 1, \dots, K$) denotes the intercept parameter for class k , $\boldsymbol{\zeta}_i$ is a q -dimensional vector of residuals assumed to follow a multivariate normal distribution $\boldsymbol{\zeta}_i \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Psi}_k)$, and $\boldsymbol{\Gamma}_k$ is a $q \times m$ parameter matrix representing the effect of $\mathbf{x}$ on the latent continuous variables $\boldsymbol{\eta}$ and assumed to be different among classes. Note that the level 2 errors $\boldsymbol{\zeta}_i$ are uncorrelated with the measurement errors $\boldsymbol{\epsilon}_i$. We may allow for class-specific effects $\boldsymbol{\Gamma}_k$ in (13) that are equal across classes.

By combining the first two levels of the GMM, we have

$$p(\mathbf{y}_i \mid \mathbf{x}_i) = \sum_{k=1}^K \pi_k \phi(\mathbf{y}_i; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k), \quad (14)$$

where $\pi_k = \Pr(C_{ik} = 1)$ is the class probability or mixing proportions satisfying $0 < \pi_k \leq 1$ and $\sum_{k=1}^K \pi_k = 1$, and $\phi(\cdot; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$ is a multivariate Gaussian density with mean $\boldsymbol{\mu}_k = \boldsymbol{\Lambda}_y(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i)$ and covariance matrix $\boldsymbol{\Sigma}_k = \boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k \boldsymbol{\Lambda}_y' + \boldsymbol{\Theta}_k$. Notice that the GMM in (14) assumes that class probability π_k is constant for each class.

At level 3 of the GMM, we assume that the class probabilities are not a constant anymore, but depend on the observed covariates. In other words, we want to know how π_k is related to an individual's background variables, such as gender and income. At this level, the categorical latent variables $\mathbf{C}_i$ represent membership of mixture components that are related to $\mathbf{x}$ through a multinomial logit regression for unordered categorical responses. Define $\pi_{ik} = \Pr(C_{ik} = 1 \mid \mathbf{x}_i)$, i.e., the probability that subject i falls into the k th class depending on the covariates $\mathbf{x}_i$. Let $\boldsymbol{\pi}_i = (\pi_{i1}, \pi_{i2}, \dots, \pi_{iK})'$ and

$$\text{logit}(\boldsymbol{\pi}_i) = \left(\log \left(\frac{\pi_{i1}}{\pi_{iK}} \right), \log \left(\frac{\pi_{i2}}{\pi_{iK}} \right), \dots, \log \left(\frac{\pi_{iK-1}}{\pi_{iK}} \right) \right)'.$$

Then,

$$\text{logit}(\boldsymbol{\pi}_i) = \boldsymbol{\alpha}_c + \boldsymbol{\Gamma}_c \mathbf{x}_i, \quad (15)$$

where $\boldsymbol{\alpha}_c$ is a $(K-1) \times 1$ parameter vector and $\boldsymbol{\Gamma}_c$ is a $(K-1) \times q$ parameter matrix. By combining these three levels of the GMM, we have

$$p(\mathbf{y}_i \mid \mathbf{x}_i) = \sum_{k=1}^K \pi_{ik} \phi(\mathbf{y}_i; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k). \quad (16)$$

Note that the model is not a finite mixture model anymore because the class probability is not constant with respect to i (cf. McLachlan and Peel, 2000).

3.2 GMM with generalized hyperbolic random effects

The conventional GMMs assume that the residuals ϵ and ζ have multivariate Gaussian distribution with zero means and within-class covariance matrices, respectively. We are interested in constructing a GMM with generalized hyperbolic distribution model errors, denoted by GHD-GMM. The generalized hyperbolic distribution can be represented as a normal mean-variance mixture, where the mixing weight has a GIG distribution. To this end, we introduce a latent continuous variable W with $W_{ik} \mid c_{ik} = 1 \sim \mathcal{I}(\omega_k, 1, \lambda_k)$. Accordingly, conditional on c_{ik} and w_{ik} , we assume that model errors ϵ_i and ζ_i are non-centered Gaussian error terms with distinct covariance matrices:

$$\epsilon_i \mid w_{ik}, c_{ik} = 1 \sim \mathcal{N}(w_i \beta_{yk}, w_{ik} \Theta_k), \quad (17)$$

$$\zeta_i \mid w_{ik}, c_{ik} = 1 \sim \mathcal{N}(w_i \beta_{\eta k}, w_{ik} \Psi_k), \quad (18)$$

where Θ_k is the diagonal covariance matrix for ϵ_i , and Ψ_k is the covariance matrix for ζ_i . The T -dimensional vector β_{yk} is a vector of skewness parameters, which we refer to as the skewness parameter for the measurement errors. The q -dimensional vector $\beta_{\eta k}$ is the vector of skew parameters for the continuous latent variables η_i . Then, based on (13) and (17), the observed random variables $\mathbf{Y}_i$, conditional on η_i , c_{ik} , and w_{ik} , follow a conditional Gaussian distribution of the form

$$\mathbf{Y}_i \mid \eta_i, w_{ik}, c_{ik} = 1 \sim \mathcal{N}(\Lambda_y \eta_i + w_i \beta_{yk}, w_{ik} \Theta_k). \quad (19)$$

And, based on (13) and (18),

$$\eta_i \mid \mathbf{x}_i, w_{ik}, c_{ik} = 1 \sim \mathcal{N}(\alpha_k + \Gamma_k \mathbf{x}_i + w_{ik} \beta_{\eta k}, w_{ik} \Psi_k). \quad (20)$$

and, from the preceding equations, we have the conditional distribution

$$\mathbf{Y}_i \mid \mathbf{x}_i, w_{ik}, c_{ik} = 1 \sim \mathcal{N}(\mu_k + w_{ik}(\Lambda_y \beta_{\eta k} + \beta_{yk}), w_{ik} \Sigma_k), \quad (21)$$

where $\mu_k = \Lambda_y(\alpha_k + \Gamma_k \mathbf{x}_i)$ and $\Sigma_k = \Lambda_y \Psi_k \Lambda_y' + \Theta_k$. Recalling the elegant result explained in Section 2.1, we obtain the conditional distributions

$$\eta_i \mid \mathbf{x}_i, c_{ik} = 1 \sim \text{GHD}_q(\lambda_k, \omega_k, \alpha_k + \Gamma_k \mathbf{x}_i, \Psi_k, \beta_{\eta k}), \quad (22)$$

$$\mathbf{Y}_i \mid \mathbf{x}_i, c_{ik} = 1 \sim \text{GHD}_T(\lambda_k, \omega_k, \mu_k, \Sigma_k, \Lambda_y \beta_{\eta k} + \beta_{yk}). \quad (23)$$

By combining the preceding setup and level 3 of the GMM from Section 3.1, we arrive at a GMM with density

$$p(\mathbf{y}_i \mid \mathbf{x}_i) = \sum_{k=1}^K \pi_{ik} f_{\text{GHDT}}(\mathbf{y}_i; \lambda_k, \omega_k, \mu_k, \Sigma_k, \Lambda_y \beta_{\eta k} + \beta_{yk}), \quad (24)$$

where $f_{\text{GHD}T}(\cdot)$ is the density of a T -dimensional random variable following a generalized hyperbolic distribution. Note that the overall skewness for $\mathbf{Y}_i$ is $\mathbf{\Lambda}_y\boldsymbol{\beta}_{\eta k} + \boldsymbol{\beta}_{yk}$. In the above setup, the dependent observed variable $\mathbf{Y}_i$, the latent growth factors $\boldsymbol{\eta}_i$, and residual variables $\boldsymbol{\epsilon}_i$ and $\boldsymbol{\zeta}_i$ all have generalized hyperbolic distributions. Note that the distribution of the covariates $\mathbf{x}_i$ is not modelled; please refer to Muthén and Asparouhov (2015) for detailed explanations.

3.3 GMM with multivariate skew-t random effects

In this section, we are interested in extending the conventional GMM to have multivariate skew-t distribution model errors, denoted by GST-GMM. As in the case for the generalized hyperbolic distribution, the multivariate skew-t distribution also has a convenient representation as a normal mean-variance mixture; this time, the weight has an inverse-gamma distribution. In analogous fashion to the GHD-GMM, a latent continuous random variable W_{ik} is first introduced, where $W_{ik} \mid c_{ik} = 1 \sim \text{IG}(\nu_k/2, \nu_k/2)$. Accordingly, we assume that $\boldsymbol{\epsilon}_i$ and $\boldsymbol{\zeta}_i$ are non-centred Gaussian error terms with their own covariance matrices as in (17) and (18), and $\mathbf{y}_i$ and $\boldsymbol{\eta}_i$ are conditionally normally distributed as in (19) and (20). From this characterization of the multivariate skew-t distribution, the following conditional distributions are obtained:

$$\boldsymbol{\eta}_i \mid \mathbf{x}_i, c_{ik} = 1 \sim \text{GST}_q(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i, \boldsymbol{\Psi}_k, \boldsymbol{\beta}_{\eta k}, \nu_k), \quad (25)$$

$$\mathbf{Y}_i \mid \mathbf{x}_i, c_{ik} = 1 \sim \text{GST}_T(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k, \boldsymbol{\beta}_{yk} + \mathbf{\Lambda}_y \boldsymbol{\beta}_{\eta k}, \nu_k), \quad (26)$$

where $\boldsymbol{\mu}_k$ and $\boldsymbol{\Sigma}_k$ are as described above and ν_k is a concentration parameter (i.e., the degrees of freedom). Similarly, we arrive at a GMM with a multivariate skew-t distribution

$$p(\mathbf{y}_i \mid \mathbf{x}_i) = \sum_{k=1}^K \pi_{ik} f_{\text{GST}T}(\mathbf{y}_i; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k, \boldsymbol{\beta}_{yk} + \mathbf{\Lambda}_y \boldsymbol{\beta}_{\eta k}, \nu_k), \quad (27)$$

where π_{ik} is defined as in (15). In this setup, the random variable $\mathbf{Y}_i$, the latent growth factors $\boldsymbol{\eta}_i$, and the residual variables $\boldsymbol{\epsilon}_i$ and $\boldsymbol{\zeta}_i$ all follow multivariate skew-t distributions.

3.4 Comments on the GHD-GMM and GST-GMM

In the preceding extensions of GMMs, recalling that the overall skewness for $\mathbf{Y}_i$ is $\mathbf{\Lambda}_y \boldsymbol{\beta}_{\eta k} + \boldsymbol{\beta}_{yk}$, so there are a total of $T + q$ skewness parameters. Hence, the skewness parameters $\boldsymbol{\beta}_{yk}$ and $\boldsymbol{\beta}_{\eta k}$ are subject to identifiability issues, because no more than T skewness parameters can be identified from the T -dimensional $\mathbf{Y}_i$. Therefore, two special formulations are considered in this paper. The first formulation is where $\boldsymbol{\beta}_{yk} = \mathbf{0}$. In this formulation, the residuals for $\mathbf{Y}_i$ or the measurement errors are not skewed, i.e., $\boldsymbol{\epsilon}_i \mid w_{ik}, c_{ik} = 1 \sim \mathcal{N}(\mathbf{0}, w_{ik} \boldsymbol{\Theta}_k)$. All of the skewness in the data is assumed to come from the distribution of latent factors. The

second special formulation is the case where $\beta_{\eta k} = \mathbf{0}$. In this formulation, the residuals for the latent factors η are symmetric, i.e., $\zeta_i \mid w_{ik}, c_{ik} = 1 \sim \mathcal{N}(\mathbf{0}, w_{ik} \Psi_k)$. Accordingly, all of the skewness in the data is assumed to come from the residuals of $\mathbf{Y}_i$ or the measurement errors. In practice, we would want as much of the skewness as possible in the observed data $\mathbf{Y}_1, \dots, \mathbf{Y}_n$ to be explained through the latent factors. There appears to be no optimal strategy with respect to which skewness parameter to estimate. Accordingly, four statistical models, differing with respect to the distributions of measurement errors and random effects for the first two levels of the GMM, are employed and compared. These models are as follows:

- **Model I:** A model with independent multivariate generalized hyperbolic random effects and measurement errors while assuming all of the skewness in the data comes from the distribution of latent factors (i.e., GHD-GMM under $\beta_{yk} = \mathbf{0}$).
- **Model II:** A model with independent multivariate generalized hyperbolic random effects and measurement errors while assuming all of the skewness in the data comes from the residuals of $\mathbf{Y}$ (i.e., GHD-GMM under $\beta_{\eta k} = \mathbf{0}$).
- **Model III:** A model with independent multivariate skew-t random effects and measurement errors while assuming all of the skewness in the data comes from the distribution of latent factors (i.e., GST-GMM under $\beta_{yk} = \mathbf{0}$).
- **Model IV:** A model with independent multivariate skew-t random effects and measurement errors while assuming all of the skewness in the data comes from the residuals of $\mathbf{Y}$ (i.e., GST-GMM under $\beta_{\eta k} = \mathbf{0}$).

Take Model I (i.e., GHD-GMM under $\beta_{yk} = \mathbf{0}$) as an example. For different trajectory classes, the parameters $\lambda_k, \omega_k, \alpha_k, \beta_{\eta k}, \Theta_k, \Psi_k$, and Γ_k may be different across classes, or may be the same across the classes. By imposing constraints on all these parameters (different or the same across classes), we obtain a family of GHD-GMM models. In this paper, we only consider two models, one model assumes that the parameters $\lambda_k, \omega_k, \alpha_k, \beta_{\eta k}, \Theta_k, \Psi_k$, and Γ_k are different across classes, we call this model as the general model. The second model assumes that only the parameter α_k is different across classes while all the other parameters are the same across classes, i.e., $\lambda_k = \lambda, \omega_k = \omega, \beta_{\eta k} = \beta_{\eta}, \Theta_k = \Theta, \Psi_k = \Psi$, and $\Gamma_k = \Gamma$ for $k = 1, 2, \dots, K$; we call this model the most constrained model.

4 Parameter Estimation

4.1 The EM algorithm for Model I

For our GHD-GMM under $\beta_{yk} = \mathbf{0}$, the complete-data comprise the observed outcome data $\mathbf{y}_1, \dots, \mathbf{y}_n$, the covariates $\mathbf{x}_1, \dots, \mathbf{x}_n$, together with the latent categorical variables $\mathbf{c}_1, \dots, \mathbf{c}_n$,

the latent growth factors $\boldsymbol{\eta}_1, \dots, \boldsymbol{\eta}_n$, and the latent w_{ik} . The observed-data log-likelihood can be expressed as follows:

$$\log \mathcal{L} = \sum_{i=1}^n \log p(\mathbf{y}_i \mid \mathbf{x}_i), \quad (28)$$

where

$$p(\mathbf{y}_i \mid \mathbf{x}_i) = \sum_{k=1}^K \pi_{ik} f_{\text{GHDT}}(\mathbf{y}_i; \lambda_k, \omega_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k, \boldsymbol{\Lambda}_y \boldsymbol{\beta}_{\eta k}) \quad (29)$$

and π_{ik} is defined as in connection with (15).

Now, $\mathbf{Y}_i \mid \mathbf{x}_i, w_{ik}, c_{ik} = 1 \sim \mathcal{N}(\boldsymbol{\mu}_k + w_{ik} \boldsymbol{\Lambda}_y \boldsymbol{\beta}_{\eta k}, w_{ik} \boldsymbol{\Sigma}_k)$ independently for $i = 1, \dots, n$, $W_{ik} \mid c_{ik} = 1 \sim \mathcal{I}(\omega_k, 1, \tilde{\lambda}_k)$, and so, from Bayes's theorem, $W_{ik} \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1 \sim \text{GIG}(\psi_k, \chi_{ik}, \tilde{\lambda}_k)$ with $\psi_k = \omega_k + \boldsymbol{\beta}'_{\eta k} \boldsymbol{\Lambda}'_y \boldsymbol{\Sigma}_k^{-1} \boldsymbol{\Lambda}_y \boldsymbol{\beta}_{\eta k}$, $\chi_{ik} = \omega_k + \delta(\mathbf{y}_i, \boldsymbol{\mu}_k \mid \boldsymbol{\Sigma}_k)$, and $\tilde{\lambda}_k = \lambda_k - T/2$. It follows that

$$\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, w_{ik}, c_{ik} = 1 \sim \mathcal{N}(\mathbf{V}_k (\boldsymbol{\Psi}_k^{-1} (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k}) + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i), w_{ik} \mathbf{V}_k), \quad (30)$$

where $\mathbf{V}_k = (\boldsymbol{\Psi}_k^{-1} + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y)^{-1}$. The result in (30) is used to estimate the latent growth factors $\boldsymbol{\eta}_i$, and a detailed proof thereof is given in Appendix A. Therefore, the complete-data likelihood is given by

$$\mathcal{L}_c(\vartheta) = \prod_{i=1}^n \prod_{k=1}^K [\pi_{ik} \phi(\mathbf{y}_i \mid \boldsymbol{\Lambda}_y \boldsymbol{\eta}_i, w_{ik} \boldsymbol{\Theta}_k) \phi(\boldsymbol{\eta}_i \mid \boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k}, w_{ik} \boldsymbol{\Psi}_k) h(w_{ik} \mid \omega_k, \lambda_k)]^{c_{ik}},$$

with the same notation used previously, where $h(w_{ik} \mid \omega_k, \lambda_k)$ is the density of a GIG distribution in (7) with $\eta = 1$.

After some algebra, the complete-data log-likelihood is given by

$$\mathcal{L}_c(\vartheta \mid \mathbf{y}, \mathbf{x}) = \mathcal{L}_{1c}(\pi) + \mathcal{L}_{2c}(\boldsymbol{\Theta}_k) + \mathcal{L}_{2c}(\boldsymbol{\alpha}_k, \boldsymbol{\beta}_{\eta k}, \boldsymbol{\Psi}_k, \boldsymbol{\Gamma}_k) + \mathcal{L}_{4c}(\boldsymbol{\lambda}, \boldsymbol{\omega}), \quad (31)$$

where $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_K)$ and $\boldsymbol{\omega} = (\omega_1, \dots, \omega_K)$, and

$$\begin{aligned} \mathcal{L}_{1c} &= \sum_{i=1}^n \sum_{k=1}^K c_{ik} \log \pi_{ik}, \\ \mathcal{L}_{2c} &= \sum_{i=1}^n \sum_{k=1}^K c_{ik} \left\{ \frac{1}{2} \log |\boldsymbol{\Theta}_k^{-1}| - \frac{1}{2w_{ik}} \mathbf{y}'_i \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i + \frac{1}{w_{ik}} \mathbf{y}'_i \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y \boldsymbol{\eta}_i - \frac{1}{2w_{ik}} \boldsymbol{\eta}'_i \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y \boldsymbol{\eta}_i \right\} + C_1, \\ \mathcal{L}_{3c} &= \sum_{i=1}^n \sum_{k=1}^K c_{ik} \left\{ \frac{1}{2} \log |\boldsymbol{\Psi}_k^{-1}| - \frac{1}{2w_{ik}} \boldsymbol{\eta}'_i \boldsymbol{\Psi}_k^{-1} \boldsymbol{\eta}_i + \frac{1}{w_{ik}} (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i)' \boldsymbol{\Psi}_k^{-1} \boldsymbol{\eta}_i + \boldsymbol{\beta}'_{\eta k} \boldsymbol{\Psi}_k^{-1} \boldsymbol{\eta}_i \right. \\ &\quad \left. - \frac{1}{2w_{ik}} (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i)' \boldsymbol{\Psi}_k^{-1} (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) - (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i)' \boldsymbol{\Psi}_k^{-1} \boldsymbol{\beta}_{\eta k} - \frac{w_{ik}}{2} \boldsymbol{\beta}'_{\eta k} \boldsymbol{\Psi}_k^{-1} \boldsymbol{\beta}_{\eta k} \right\} + C_2, \\ \mathcal{L}_{4c} &= \sum_{i=1}^n \sum_{k=1}^K c_{ik} \left\{ (\lambda_k - 1) \log w_{ik} - \log K_{\lambda_k}(\omega_k) - \frac{\omega_k}{2} \left(w_{ik} + \frac{1}{w_{ik}} \right) \right\}, \end{aligned}$$

where C_1 and C_2 are constants with respect to model parameters.

In the E-step, we compute the conditional expectation of $\mathcal{L}_c(\boldsymbol{\vartheta} \mid \mathbf{y}, \mathbf{x})$ given in (32), denoted $\mathcal{Q}$. First, let p_{ik} denote the probability that the i th observation belongs to the k th component of the mixture, and is updated by

$$p_{ik} := \mathbb{E}[C_{ik} \mid \mathbf{y}_i, \mathbf{x}_i] = \frac{\pi_{ik} f_{\text{GHDT}}(\mathbf{y}_i; \lambda_k, \omega_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k, \boldsymbol{\Lambda}_y \boldsymbol{\beta}_{\eta k})}{\sum_{l=1}^K \pi_{il} f_{\text{GHDT}}(\mathbf{y}_i; \lambda_l, \omega_l, \boldsymbol{\mu}_l, \boldsymbol{\Sigma}_l, \boldsymbol{\Lambda}_y \boldsymbol{\beta}_{\eta l})}.$$

The resulting E-step quantiles regarding the latent growth factors $\boldsymbol{\eta}$ and the latent continuous variable W are given by

$$\begin{aligned} E_{1ik} &:= \mathbb{E}[W_i \mid \mathbf{x}_i, \mathbf{y}_i, c_{ik} = 1] = \sqrt{\frac{\chi_{ik}}{\psi_k}} \frac{K_{\tilde{\lambda}_k+1}(\sqrt{\psi_k \chi_{ik}})}{K_{\tilde{\lambda}_k}(\sqrt{\psi_k \chi_{ik}})}, \\ E_{2ik} &:= \mathbb{E}[1/W_i \mid \mathbf{x}_i, \mathbf{y}_i, c_{ik} = 1] = \sqrt{\frac{\psi_k}{\chi_{ik}}} \frac{K_{\tilde{\lambda}_k+1}(\sqrt{\psi_k \chi_{ik}})}{K_{\tilde{\lambda}_k}(\sqrt{\psi_k \chi_{ik}})} - \frac{2\tilde{\lambda}_k}{\chi_{ik}}, \\ E_{3ik} &:= \mathbb{E}[\log W_i \mid \mathbf{x}_i, \mathbf{y}_i, c_{ik} = 1] = \log \left(\sqrt{\frac{\chi_{ik}}{\psi_k}} \right) + \frac{1}{K_{\tilde{\lambda}_k}(\sqrt{\psi_k \chi_{ik}})} \frac{\partial}{\partial \tilde{\lambda}_k} K_{\tilde{\lambda}_k}(\sqrt{\psi_k \chi_{ik}}), \\ E_{4ik} &:= \mathbb{E}[\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1] = \mathbf{V}_k (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + E_{1ik} \boldsymbol{\beta}_{\eta k}) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i), \\ E_{5ik} &:= \mathbb{E}[(1/W_{ik}) \boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1] = E_{2ik} \mathbf{V}_k (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i) + \mathbf{V}_k \boldsymbol{\Psi}_k^{-1} \boldsymbol{\beta}_{\eta k}, \\ E_{6ik} &:= \mathbb{E}[(1/W_{ik}) \boldsymbol{\eta}_i \boldsymbol{\eta}_i' \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1] = \mathbf{V}_k + \mathbf{V}_k (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i) \boldsymbol{\beta}_{\eta k} \boldsymbol{\Psi}_k^{-1} \mathbf{V}_k \\ &\quad + E_{2ik} \mathbf{V}_k (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i) (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i)' \mathbf{V}_k \\ &\quad + \mathbf{V}_k \boldsymbol{\Psi}_k^{-1} \boldsymbol{\beta}_{\eta k}' (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i)' \mathbf{V}_k + E_{1ik} \mathbf{V}_k \boldsymbol{\Psi}_k^{-1} \boldsymbol{\beta}_{\eta k} \boldsymbol{\beta}_{\eta k}' \boldsymbol{\Psi}_k^{-1} \mathbf{V}_k, \end{aligned}$$

and ψ_k , χ_{ik} , and $\tilde{\lambda}_k$ are as previously defined. These attractive closed forms for E_{1ik} , E_{2ik} , and E_{3ik} exist because $W_{ik} \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1 \sim \text{GIG}(\psi_k, \chi_{ik}, \tilde{\lambda}_k)$, and so we can use the formulae as in (8). The existence of these attractive closed forms for E_{4ik} , E_{5ik} , and E_{6ik} is due to the conditional Gaussian distribution of $\boldsymbol{\eta}$ as in (30).

In the M-step, we maximize $\mathcal{Q}$ with respect to the model parameters to get the updates. With respect to the parameters $\boldsymbol{\alpha}_c$ and $\boldsymbol{\Gamma}_c$, the M-step maximizes

$$\sum_{i=1}^n \sum_{k=1}^K p_{ik} \log \pi_{ik}, \quad (32)$$

which may be viewed as a multinomial logistic regression with fractional observations p_{ik} . The parameters ω_k and λ_k are estimated by maximizing the following function

$$q_k(\omega_k, \lambda_k) = -\log K_{\lambda_k}(\omega_k) + (\lambda_k - 1) \bar{d}_k - \frac{\omega_k}{2} (\bar{a}_k + \bar{b}_k), \quad (33)$$

where $n_k = \sum_{i=1}^n p_{ik}$, $\bar{a}_k = \frac{1}{n_k} \sum_{i=1}^n p_{ik} E_{1ik}$, $\bar{b}_k = \frac{1}{n_k} \sum_{i=1}^n p_{ik} E_{2ik}$, and $\bar{d}_k = \frac{1}{n_k} \sum_{i=1}^n p_{ik} E_{3ik}$. The associated updates are

$$\begin{aligned}\hat{\lambda}_k &= \bar{c}_k \hat{\lambda}_k^{\text{prev}} \left[\frac{\partial}{\partial t} \log K_t(\hat{\omega}_k^{\text{prev}}) \Big|_{t=\hat{\lambda}_k^{\text{prev}}} \right]^{-1}, \\ \hat{\omega}_k &= \hat{\omega}_k^{\text{prev}} - \left[\frac{\partial}{\partial t} q_k(t, \hat{\lambda}_k) \Big|_{t=\hat{\omega}_k^{\text{prev}}} \right] \left[\frac{\partial^2}{\partial t^2} q_k(t, \hat{\lambda}_k) \Big|_{t=\hat{\omega}_k^{\text{prev}}} \right]^{-1},\end{aligned}$$

where the superscript “prev” means the previous estimate — refer to Browne and McNicholas (2015) for further details. Finally, we get the updates of the other parameters in the model:

$$\begin{aligned}\hat{\Theta}_k &= \text{diag} \left(\frac{\sum_{i=1}^n p_{ik} (E_{2ik} \mathbf{y}_i \mathbf{y}_i' - \mathbf{y}_i E_{5ik}' \Lambda_y' - \Lambda_y E_{5ik} \mathbf{y}_i' + \Lambda_y E_{6ik} \Lambda_y')}{n_k} \right), \\ \hat{\Gamma}_k &= \sum_{i=1}^n p_{ik} \left(E_{5ik} \mathbf{x}_i' - E_{2ik} \hat{\alpha}_k \mathbf{x}_i' - \hat{\beta}_k \mathbf{x}_i' \right) \left(\sum_{i=1}^n \sum_{k=1}^K p_{ik} E_{2ik} \mathbf{x}_i \mathbf{x}_i' \right)^{-1}, \\ \hat{\alpha}_k &= \frac{\bar{a}_k \sum_{i=1}^n p_{ik} (E_{5ik} - E_{2ik} \hat{\Gamma}_k \mathbf{x}_i) - \sum_{i=1}^n p_{ik} E_{4ik} + \sum_{i=1}^n p_{ik} \hat{\Gamma}_k \mathbf{x}_i}{n_k (\bar{a}_k \bar{b}_k - 1)}, \\ \hat{\beta}_{\eta k} &= \frac{\bar{b}_k \sum_{i=1}^n p_{ik} (E_{4ik} - \hat{\Gamma}_k \mathbf{x}_i) - \sum_{i=1}^n p_{ik} E_{5ik} + \sum_{i=1}^n p_{ik} E_{2ik} \hat{\Gamma}_k \mathbf{x}_i}{n_k (\bar{b}_k \bar{b}_k - 1)}, \\ \hat{\Psi}_k &= \frac{1}{n_k} \sum_{i=1}^n p_{ik} \left(E_{6ik} - \hat{\beta}_k E_{4ik}' - E_{5ik} (\hat{\alpha}_k + \hat{\Gamma}_k \mathbf{x}_i)' - E_{4ik} \hat{\beta}_k' - (\hat{\alpha}_k + \hat{\Gamma}_k \mathbf{x}_i) E_{5ik}' \right. \\ &\quad \left. + E_{2ik} (\hat{\alpha}_k + \hat{\Gamma}_k \mathbf{x}_i) (\hat{\alpha}_k + \hat{\Gamma}_k \mathbf{x}_i)' + (\hat{\alpha}_k + \hat{\Gamma}_k \mathbf{x}_i) \hat{\beta}_{\eta k}' + \hat{\beta}_{\eta k} (\hat{\alpha}_k + \hat{\Gamma}_k \mathbf{x}_i)' + E_{1ik} \hat{\beta}_{\eta k} \hat{\beta}_{\eta k}' \right).\end{aligned}$$

4.2 The EM algorithm for Model III

Similarly, for our Model III, parameter estimation is carried out within the EM algorithm framework. Suppose we observe the outcome $\mathbf{y}_i$ and the covariates $\mathbf{x}_i$ from a GMM with skew-t random effects as in (27) but with $\beta_{yk} = \mathbf{0}$. There are three sources of unobserved data: the latent categorical variables $\mathbf{c}_i$, the latent growth factors $\boldsymbol{\eta}_i$, and the latent w_{ik} . The complete-data log-likelihood can be expressed as follows:

$$\begin{aligned}\mathcal{L}_c(\vartheta) &= \sum_{i=1}^n \sum_{k=1}^K \pi_{ik} \left[\log \pi_{ik} + \log \phi(\mathbf{y}_i \mid \Lambda_y \boldsymbol{\eta}_i, w_{ik} \boldsymbol{\Theta}_k) \right. \\ &\quad \left. + \log \phi(\boldsymbol{\eta}_i \mid \boldsymbol{\alpha}_k + \Gamma_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k}, w_{ik} \boldsymbol{\Psi}_k) + \log f(w_{ik} \mid \nu_k/2, \nu_k/2) \right],\end{aligned}$$

where π_{ik} is defined as above in (15) and $f(w_{ik} \mid \nu_k/2, \nu_k/2)$ is the density of the inverse Gamma distribution.

The E-step requires the computation of the expected value of the complete-data log-likelihood. Note that $W_{ik} \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1 \sim \text{GIG}(\psi_k^*, \chi_{ik}^*, \lambda_k^*)$ with $\psi_k^* = \beta_{\eta k}' \Lambda_y' \Sigma^{-1} \Lambda_y \beta_{\eta k}$,

$\chi_{ik}^* = \nu_k + \delta(\mathbf{y}_i, \boldsymbol{\mu}_k \mid \boldsymbol{\Sigma}_k)$, and $\lambda_k^* = -(\lambda_k + T)/2$. So, we have convenient forms for the following expected values:

$$\begin{aligned} E_{1ik}^* &:= \mathbb{E}[W_i \mid \mathbf{x}_i, \mathbf{y}_i, c_{ik} = 1] = \sqrt{\frac{\chi_{ik}^*}{\psi_k^*}} \frac{K_{\lambda_k^*+1}(\sqrt{\psi_k^* \chi_{ik}^*})}{K_{\lambda_k^*}(\sqrt{\psi_k^* \chi_{ik}^*})}, \\ E_{2ik}^* &:= \mathbb{E}[1/W_i \mid \mathbf{x}_i, \mathbf{y}_i, c_{ik} = 1] = \sqrt{\frac{\psi_k^*}{\chi_{ik}^*}} \frac{K_{\lambda_k^*+1}(\sqrt{\psi_k^* \chi_{ik}^*})}{K_{\lambda_k^*}(\sqrt{\psi_k^* \chi_{ik}^*})} - \frac{2\lambda_k^*}{\chi_{ik}^*}, \\ E_{3ik}^* &:= \mathbb{E}[\log W_i \mid \mathbf{x}_i, \mathbf{y}_i, c_{ik} = 1] = \log \left(\sqrt{\frac{\chi_{ik}^*}{\psi_k^*}} \right) + \frac{1}{K_{\lambda_k^*}(\sqrt{\psi_k^* \chi_{ik}^*})} \frac{\partial}{\partial \lambda_k^*} K_{\lambda_k^*}(\sqrt{\psi_k^* \chi_{ik}^*}). \end{aligned}$$

We also need the expected value of the class membership, i.e.,

$$\tau_{ik} := \mathbb{E}[C_{ik} \mid \mathbf{y}_i, \mathbf{x}_i] = \frac{\pi_{ik} f_{\text{GHD}}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k, \boldsymbol{\Lambda}_y \boldsymbol{\beta}_{\eta k}, v_k)}{\sum_{l=1}^K \pi_{il} f_{\text{GHD}}(\boldsymbol{\mu}_l, \boldsymbol{\Sigma}_l, \boldsymbol{\Lambda}_y \boldsymbol{\beta}_{\eta l}, v_l)},$$

as well as the following conditional expectations, which are similar to those derived in the E-step of parameter estimation for the GHD-GMM:

$$\begin{aligned} E_{4ik}^* &:= \mathbb{E}[\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1] = \mathbf{V}_k (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + E_{1ik}^* \boldsymbol{\beta}_{\eta k}) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i), \\ E_{5ik}^* &:= \mathbb{E}[(1/W_{ik}) \boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1] = E_{2ik}^* \mathbf{V}_k (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i) + \mathbf{V}_k \boldsymbol{\Psi}_k^{-1} \boldsymbol{\beta}_{\eta k}, \\ E_{6ik}^* &:= \mathbb{E}[(1/W_{ik}) \boldsymbol{\eta}_i \boldsymbol{\eta}_i' \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1] = \mathbf{V}_k + \mathbf{V}_k (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i) \boldsymbol{\beta}_{\eta k} \boldsymbol{\Psi}_k^{-1} \mathbf{V}_k \\ &\quad + E_{2ik}^* \mathbf{V}_k (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i) (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i)' \mathbf{V}_k, \\ &\quad + \mathbf{V}_k \boldsymbol{\Psi}_k^{-1} \boldsymbol{\beta}_{\eta k}' (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i)' \mathbf{V}_k + E_{1ik}^* \mathbf{V}_k \boldsymbol{\Psi}_k^{-1} \boldsymbol{\beta}_{\eta k} \boldsymbol{\beta}_{\eta k}' \boldsymbol{\Psi}_k^{-1} \mathbf{V}_k. \end{aligned}$$

The M-step requires the computation of the parameter updates to maximize the expected value of the complete-data log-likelihood. In this step, the parameter updates for $\boldsymbol{\alpha}_c, \boldsymbol{\Gamma}_c, \boldsymbol{\alpha}_k, \boldsymbol{\beta}_{\eta k}, \boldsymbol{\Theta}_k, \boldsymbol{\Psi}_k$ and $\boldsymbol{\Gamma}_k$ are obtained in closed form and are similar to those derived in the M-step of parameter estimation for the GHD-GMM, hence are omitted here. We solve the equation

$$\log \left(\frac{\nu_k}{2} \right) + 1 - \varphi \left(\frac{\nu_k}{2} \right) - \frac{1}{n_k} \sum_{i=1}^n \tau_{ik} (E_{3ik}^* + E_{2ik}^*) = 0 \quad (34)$$

for ν_k , numerically, where $n_k = \sum_{i=1}^n \tau_{ik}$. Parameter estimation for Models II and IV is outlined in Appendix B.

5 Illustrations

5.1 Performance assessment

Although all of our illustrations are treated as genuine clustering analysis, i.e., no prior knowledge of labels is assumed, the true labels are known in each case and can be used to

evaluate the performance of our GHD-GMM and GST-GMM. We use the adjusted Rand index (ARI; Hubert and Arabie, 1985) to assess classification performance. The ARI indicates the pairwise agreement between true and predicted group memberships and, in doing so, accounts for the fact that random classification would classify some observations correctly by chance. An ARI value of 1 indicates perfect classification, a value of 0 would be expected under random classification, and a negative ARI value indicates classification that is worse than one would expect under random classification. For further details and discussion on the ARI, see Steinley (2004).

5.2 Alcoholic consumption data from the National Longitudinal Survey of Youth

The National Longitudinal Survey of Youth (NLSY) is a longitudinal study conducted by the United States Bureau of Labor Statistics with the goal of understanding the interaction between labor force participation, education, and health behaviors in children and adolescents. The sample for this study was a cohort of children who were between the ages of 12 and 17 when first interviewed in 1997. The data of interest were gathered each year between 1997 and 2011 and again in 2013 (15 total possible interviews). Each respondent provided a number between 0 and 98 that represents the number of alcoholic drinks they typically consume on a given day on which they are drinking. Tracking heterogeneity in trajectories of alcohol consumption has been a frequent topic in the social and biomedical sciences with many objectives: to determine the effect of excessive drinking over time, design interventions that can prevent excessive drinking, or model the interplay between consumption of different drugs. Because we are interested in modeling drinking behaviour over the life span, the data are shifted from representing year of interview to age. To minimize the effect of missing data, individuals who were interviewed between the ages of 16 and 19 were used for the following analyses.

We implement the Gaussian GMM via `Mplus` Version 7.1 (Muthén and Muthén, 2012). Our proposed GHD-GMM and GST-GMM are implemented in `R` and run with K ranging from 1 to 10 until the best model is obtained under each scenario. Table 1 shows the results of fitting all of the models as aforementioned for a varying number of latent classes. The BIC values show that more than eight classes are needed with the conventional GMM, two are needed with constrained Models I and IV, and three are needed for all of the other models. The BIC values for the GST-GMM and GHD-GMM are better than the BIC for the normal GMM. Notably, the BIC values for the GHD-GMM do not always improve on that for the GST-GMM. Among all fitted models, the three-cluster general GST-GMM under $\beta_{yk} = \mathbf{0}$ (i.e., general Model III) is preferable according to the BIC. It is worth mentioning that, even though the skew-t distribution is a special case of the generalized hyperbolic distribution, the GST-GMM seems to be useful in addition to the GHD-GMM.

The best-fitting model, the three-cluster skew-t, breaks the data into three groups. The first group, comprising 56% of the population, begins with low-moderate drinking (< 1 drink

Table 1: Results of fitting normal, GST, and GHD GMMs for consumption data from the National Longitudinal Survey of Youth.

GMM-Normal (constrained)				GMM-Normal (general)		
Classes	Log-likelihood	Free paras	BIC	Log-likelihood	Free paras	BIC
1	-14983.42	9	-30030.27	-14983.42	9	-30030.27
2	-14623.41	12	-29331.40	-12671.88	19	-25477.69
3	-14330.00	15	-28765.72	-12233.95	29	-24672.31
4	-14182.66	18	-28492.19	-12119.21	39	-24513.30
5	-14076.42	21	-28300.85	-12027.60	49	-24400.58
6	-14015.58	24	-28200.32	-11950.97	59	-24317.78
7	-13980.78	27	-28151.86	-11906.09	69	-24298.53
8	-13937.17	30	-28085.80	-11870.44	79	-24297.69
9	-13916.40	33	-28065.38			
GHD-GMM (Model I, constrained)				GHD-GMM (Model I, general)		
Classes	Log-likelihood	Free paras	BIC	Log-likelihood	Free paras	BIC
1	-12403.20	13	-24898.19	-12403.28	13	-24898.19
2	-12315.75	16	-24744.27	-12119.53	27	-24429.36
3	-12315.50	19	-24764.91	-11958.92	41	-24206.82
4				-11953.84	55	-24295.33
GHD-GMM (Model II, constrained)				GHD-GMM (Model II, general)		
Classes	Log-likelihood	Free paras	BIC	Log-likelihood	Free paras	BIC
1	-12399.68	15	-24883.94	-12399.68	15	-24905.09
2	-12312.27	18	-24737.32	-12166.47	31	-24551.45
3	-12288.26	21	-24717.49	-12002.12	47	-24335.51
4	-12287.98	24	-24745.12	-11956.47	63	-24356.99
GST-GMM (Model III, constrained)				GST-GMM (Model III, general)		
Classes	Log-likelihood	Free paras	BIC	Log-likelihood	Free paras	BIC
1	-12421.85	12	-24928.28	-12421.92	12	-24928.42
2	-12352.31	15	-24810.34	-12151.6	25	-24479.41
3	-12340.61	18	-24808.10	-11966.84	38	-24201.52
4	-12348.28	21	-24844.58	-11925.67	51	-24210.82
GST-GMM (Model IV, constrained)				GST-GMM (Model IV, general)		
Classes	Log-likelihood	Free paras	BIC	Log-likelihood	Free paras	BIC
1	-12418.18	14	-24935.05	-12418.19	14	-24935.05
2	-12348.01	17	-24748.06	-12118.12	29	-24440.64
3	-12347.48	20	-24756.18	-11990.60	44	-24291.33
4				-11938.08	59	-24292.00

per drinking day), slightly increases during adolescence, and by age 19 the average drinks per drinking day is at about 1. These can be considered “consistent low” drinkers. Although the intercept for this class is heavily positively skewed (intercept skewness = 2.59), the slope is not (intercept skewness = 0.03), which indicates that the individual slopes are nearly

normally distributed around the class slope of 0.21. The second class, comprising 23% of the population, are what will be called the “decreasing” drinkers. This class has an intercept of around five drinks per drinking day (a drinking binge) and ends at about 3 drinks per drinking day (just below the amount considered a drinking binge).¹ The intercept is again positively skewed (intercept skewness = 2.90) but the slope is negatively skewed (slope skewness = -0.78), suggesting that individuals in this class decrease their consumption quickly over the period of adolescence. The third class, comprising 20% of the population, will be called the “increasing moderate” drinkers. Their initial level of drinking is around 2.87 drinks per drinking day (less than a binge) and this increases during adolescence, ending at age 19 around 7 drinks per drinking day (far above a drinking binge). Both the slope and intercept are slightly positively skewed (intercept skewness = 0.48, slope skewness = 0.41).

These results suggest that, during adolescence, which is typically a time when alcohol consumption is initiated, individuals will have different reactions to the exposure to alcohol given their previous experience – those individuals who are low drinkers will tend to continue to be low drinkers, those who have already consumed alcohol heavily will begin to taper back to safe levels (alluding to these individuals “knowing their limits” when it comes to alcohol), and those who are only at moderate levels tend to increase to heavy drinking. This model can be useful because it indicates which 15-year-olds should be the target of interventions if the goal is to prevent heavy drinking in late adolescence. Although the high drinkers may appear to be the most likely to develop problems related to alcohol, they may “grow out” of their alcohol consumption; therefore, it may be better to focus efforts on the 15-year-olds that only drink at moderate levels.

Other models tend to find a similar developmental pattern. For instance, the three-class GHD model (which has a very similar BIC to the three-class skew-t model) demonstrates a similar pattern in the clusters: one consistent low, one high but decreasing, and one moderate and increasing. This suggests that the same pattern endures regardless of the distributional assumptions. However, the cluster proportions differ slightly (58%, 24%, and 17%, for the low, high/decreasing, and moderate/increasing classes, respectively), which seems to suggest that the GHD model classifies more individuals into the “low” class than the skew-t model. If the goal of the analysis is to identify groups to target for interventions for the prevention of alcoholism, the proportions found in the skew-t model might be preferred as they create population groups that are larger. Therefore, interventions targeting this group may have a greater impact on the population than those targeting a smaller group.

¹The World Health Organization defines heavy episodic drinking (also called a drinking “binge”) as the consumption of 60 or more grams of alcohol on one occasion (www.who.int/gho/alcohol/consumption_patterns/heavy_episodic_drinkers_text/en/), which is about four standard drinks (www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/what-standard-drink).

5.3 Simulation Studies

In addition to the real data application of our proposed model, simulation studies are carried out to further illustrate the performance of the proposed GST-GMM and GHD-GMM models in recovering the underlying model parameters and the clustering ability. We use the relationship between the generalized hyperbolic distribution and the Gaussian distribution (cf. Section 2) to generate GHD-GMM and GST-GMM data. Data are generated in a number of scenarios: linear and quadratic GMMs with different distributions of the measurement errors and random effects, resulting in four distinct simulation experiments. We analyzed the data in several different ways: using a normal distribution via `Mplus`, using the proposed constrained and general GST-GMM and GHD-GMM models under $\beta_{yk} = \mathbf{0}$, and using the proposed constrained and general GST-GMM and GHD-GMM models under $\beta_{\eta k} = \mathbf{0}$. The purpose of analyzing the simulated data in this way is to compare our proposed GHD-GMM and GST-GMM models with the Gaussian GMM developed by Muthén and colleagues, which dominates the literature on GMMs.

In the first simulation experiment, the dataset is generated by a two-class general GHD-GMM under $\beta_{\eta k}$ with linear growth and five time points ($n_1 = n_2 = 400$). In the second simulation experiment, the dataset is generated by a two-class general GHD-GMM under β_{yk} with quadratic growth and thirty time points ($n_1 = n_2 = 200$). In the third simulation experiment, the simulated dataset is generated by a three-class general GST-GMM under $\beta_{\eta k}$ with linear growth and eight time points ($n_1 = n_2 = n_3 = 500$). In the fourth simulation experiment, the dataset is generated by a two-class general GST-GMM under β_{yk} with quadratic growth and twenty time points ($n_1 = n_2 = 500$). Individual trajectories for these four simulation experiments are plotted in Figure 1.

First, we evaluate the ability of our proposed model to recover underlying parameters. To this end, 100 datasets are generated for each of the four simulation experiments. True values and the means of the parameter estimates with their associated standard deviations are summarized in Tables 2–5. The results for each of the four simulation experiments show that the means of parameter estimates are close to the true values with small standard deviations; hence, our proposed approach is effective at parameter recovery.

Second, we compare the two formulations for extending GMMs. One hundred datasets are generated for the four simulation experiments above and analyzed using GMMs developed herein. The means of the BIC, the ARI, and the misclassification rates (ERR) are summarized in Table 6. As anticipated, the best models obtained are those with underlying true data structures. There is, however, one disadvantage with Model II and Model IV. In terms of model complexity, Model II has $K(T - q)$ more parameters than Model I, and Model IV has $K(T - q)$ more parameters than Model III. Hence, Models II and IV need larger sample sizes as small class sizes can create problems, such as singularity of the covariance matrix and slow or non-convergence of the EM algorithm.

Finally, we compare our proposed Models I and III with the Gaussian GMMs (via `Mplus`). Herein, 100 datasets are generated, as described before, for the second simulation experi-




Figure 1: Individual observation trajectories plots for the four simulation experiments.

Table 2: Key model parameters as well as means and standard deviations of the associated parameter estimations from the 100 runs for the first simulation experiment.

	True values	Means	Standard deviations
α_1	$(4, 5)'$	$(4.00, 5.00)'$	$(0.11, 0.08)'$
α_2	$(2, 3)'$	$(1.99, 2.98)'$	$(0.18, 0.11)'$
β_1	$(1, -1, 1, 1, 1)'$	$(0.94, -0.92, 0.93, 0.94, 0.93)'$	$(0.35, 0.33, 0.39, 0.44, 0.51)'$
β_2	$(-1, 1, -1, -1, -1)'$	$(-0.77, 0.83, -0.72, -0.68, -0.67)'$	$(0.27, 0.36, 0.33, 0.43, 0.55)'$
λ_1	-1	-0.7	0.73
λ_2	-2	-1.05	0.85
ω_1	2	2.02	0.32
ω_2	3	3.11	0.58
Ψ_1	$\begin{bmatrix} 1 & 0 \\ 0 & 0.7 \end{bmatrix}$	$\begin{bmatrix} 0.92 & 0.00 \\ 0.00 & 0.65 \end{bmatrix}$	$\begin{bmatrix} 0.34 & 0.08 \\ 0.08 & 0.22 \end{bmatrix}$
Ψ_2	$\begin{bmatrix} 1.5 & 0 \\ 0 & 0.8 \end{bmatrix}$	$\begin{bmatrix} 1.20 & -0.01 \\ -0.01 & 0.63 \end{bmatrix}$	$\begin{bmatrix} 0.36 & 0.08 \\ 0.08 & 0.18 \end{bmatrix}$

ment and analyzed with different distributions: the generalized hyperbolic distribution via our proposed Model I, the multivariate skew-t distribution via our proposed Model III, and Gaussian GMMs via the **Mplus** software. Table 7 summarizes the percentage of the replications favoured by the BIC when analyzing those 100 generated datasets for 1–6 latent

Table 3: Key model parameters as well as means and standard deviations of the associated parameter estimations from the 100 runs for the second simulation experiment.

	True values	Means	Standard deviations
α_1	$(15, 8, -6)'$	$(14.98, 8.00, -6.00)'$	$(0.10, 0.09, 0.15)'$
α_2	$(-14, -10, 6)'$	$(-14.06, -9.96, 6.02)'$	$(0.24, 0.20, 0.16)'$
β_1	$(1, 1, 1)'$	$(1.05, 1.01, 1)'$	$(0.35, 0.32, 0.39)'$
β_2	$(-1, -1, -1)'$	$(-0.89, -0.95, -0.95)'$	$(0.24, 0.25, 0.27)'$
λ_1	-1	-0.52	0.70
λ_2	2	1.05	1.28
ω_1	2	2.02	0.31
ω_2	3	2.92	0.51
Ψ_1	$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0.7 & 0 \\ 0 & 0 & 2 \end{bmatrix}$	$\begin{bmatrix} 0.99 & 0.00 & 0.00 \\ 0.00 & 0.70 & 0.00 \\ 0.00 & 0.00 & 1.99 \end{bmatrix}$	$\begin{bmatrix} 0.33 & 0.07 & 0.12 \\ 0.07 & 0.21 & 0.11 \\ 0.12 & 0.11 & 0.62 \end{bmatrix}$
Ψ_2	$\begin{bmatrix} 1.5 & 0 & 0 \\ 0 & 0.8 & 0 \\ 0 & 0 & 0.9 \end{bmatrix}$	$\begin{bmatrix} 1.38 & 0.00 & 0.01 \\ 0.00 & 0.74 & -0.01 \\ 0.01 & -0.01 & 0.84 \end{bmatrix}$	$\begin{bmatrix} 0.39 & 0.08 & 0.08 \\ 0.08 & 0.21 & 0.08 \\ 0.08 & 0.08 & 0.21 \end{bmatrix}$

Table 4: Key model parameters as well as means and standard deviations of the associated parameter estimations from the 100 runs for the third simulation experiment.

	True values	Means	Standard deviations
α_1	$(4, 5)'$	$(4.01, 4.98)'$	$(0.11, 0.10)'$
α_2	$(0, 0)'$	$(-0.01, 0.01)'$	$(0.07, 0.08)'$
α_3	$(-4, -5)'$	$(-3.91, -4.9)'$	$(0.81, 1.01)'$
β_1	$(1, 1)'$	$(1.00, 1.01)'$	$(0.10, 0.09)'$
β_2	$(0, 0)'$	$(-0.01, -0.02)'$	$(0.17, 0.19)'$
β_3	$(-1, -1)'$	$(-0.99, -0.98)$	$(0.24, 0.22)'$
ν_1	7	7.09	0.61
ν_2	5	4.97	0.41
ν_3	6	6.08	0.50
Ψ_1	$\begin{bmatrix} 1 & 0 \\ 0 & 0.7 \end{bmatrix}$	$\begin{bmatrix} 1.00 & 0.01 \\ 0.01 & 0.68 \end{bmatrix}$	$\begin{bmatrix} 0.07 & 0.05 \\ 0.05 & 0.07 \end{bmatrix}$
Ψ_2	$\begin{bmatrix} 0.7 & 0 \\ 0 & 0.6 \end{bmatrix}$	$\begin{bmatrix} 0.72 & 0.03 \\ 0.03 & 0.64 \end{bmatrix}$	$\begin{bmatrix} 0.28 & 0.32 \\ 0.32 & 0.40 \end{bmatrix}$
Ψ_3	$\begin{bmatrix} 1.5 & 0 \\ 0 & 0.8 \end{bmatrix}$	$\begin{bmatrix} 1.36 & 0.00 \\ 0.00 & 0.76 \end{bmatrix}$	$\begin{bmatrix} 0.27 & 0.07 \\ 0.07 & 0.08 \end{bmatrix}$

classes (note that 6 latent classes were never selected, cf. Table 7). It is not surprising that the Gaussian GMMs overestimate the number of classes, pointing to five classes 33% of the time and to four classes 67% of the time. Because the true data are generated from the generalized hyperbolic distribution, the Gaussian GMMs need to extract more latent classes

Table 5: Key model parameters as well as means and standard deviations of the associated parameter estimations from the 100 runs for the fourth simulation experiment.

	True values	Means	Standard deviations
α_1	$(8, 7, -3)'$	$(7.95, 7.01, -2.93)'$	$(0.18, 0.11, 0.05)'$
α_2	$(-8, -7, 3)'$	$(-7.98, -6.99, 2.93)'$	$(0.21, 0.12, 0.05)'$
β_1	$(1, 1, 1, 1, 1, 1, 1)'$	$(1.04, 0.98, 0.78, 0.44, -0.02, -0.62, -1.35, -2.21)'$	$(0.16, 0.17, 0.18, 0.18, 0.18, 0.18, 0.22, 0.32)'$
β_2	$(-1, -1, -1, -1, -1, -1, -1, -1)'$	$(-1.02, -0.96, -0.78, -0.46, -0.01, 0.56, 1.26, 2.09)'$	$(0.17, 0.15, 0.16, 0.16, 0.18, 0.22, 0.29, 0.40)'$
ν_1	7	7.43	0.80
ν_2	6	6.27	0.59
Ψ_1	$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0.7 & 0 \\ 0 & 0 & 0.8 \end{bmatrix}$	$\begin{bmatrix} 0.99 & 0.00 & 0.01 \\ 0.00 & 0.71 & 0.00 \\ 0.01 & 0.00 & 0.80 \end{bmatrix}$	$\begin{bmatrix} 0.11 & 0.07 & 0.05 \\ 0.07 & 0.06 & 0.04 \\ 0.05 & 0.04 & 0.06 \end{bmatrix}$
Ψ_2	$\begin{bmatrix} 1.5 & 0 & 0 \\ 0 & 0.8 & 0 \\ 0 & 0 & 0.9 \end{bmatrix}$	$\begin{bmatrix} 1.49 & 0.01 & 0.01 \\ 0.01 & 0.79 & 0.01 \\ 0.01 & 0.01 & 0.90 \end{bmatrix}$	$\begin{bmatrix} 0.20 & 0.10 & 0.08 \\ 0.10 & 0.09 & 0.04 \\ 0.08 & 0.04 & 0.07 \end{bmatrix}$

Table 6: Comparison of results — including average BIC, ARI, and EER values — for Models I–IV.

	Model I				Model II			
	Free paras	BIC	ARI	ERR	Free paras	BIC	ARI	ERR
1st simulation (model II)	29	-16018.99	0.92	0.02	35	-15366.59	0.97	0.01
2nd simulation (model I)	89	-69895.48	1.00	0.00	143	-70507.17	1.00	0.00
	Model IV				Model III			
	Free paras	BIC	ARI	ERR	Free paras	BIC	ARI	ERR
3rd simulation (model IV)	50	-101669.80	1.00	0.00	68	-101587.60	1.00	0.00
4th simulation (model III)	67	-37271.48	1.00	0.00	101	-37265.55	1.00	0.00

to capture the observed variable distribution. The generalized hyperbolic and skew-t GMMs (i.e., the proposed Models I and III) correctly choose the number of classes 100% of the time. It is noteworthy to mention that the best model, based on the BIC, is a two-class generalized hyperbolic GMM consistently. Typical analysis results for fitting the Gaussian, the multi-variate skew-t, and the generalized hyperbolic GMMs to the second simulation experiment are given in Table 8.

Table 7: Percent preferred by the BIC when analyzing the second simulation experiment with Model I, Model III, and GMM along with number of classes.

	Number of classes				
	1	2	3	4	5
Model I	0	100	0	0	0
Model III	0	100	0	0	0
GMM	0	0	0	67	33

Table 8: Results of fitting the Gaussian, GST, and GHD to the second simulation experiment.

Model	K	Free paras	Log-likelihood	BIC	ARI	ERR
Model I	1	64	-35261.30	-70906.06	0	0.50
	2	129	-34283.40	-69339.69	1	0
	3	194	-34231.13	-69624.60	0.86	0.08
Model III	1	63	-35295.15	-70967.76	0	0.5
	2	127	-34292.80	-69346.20	1	0
	3	191	-34240.11	-69624.59	0.86	0.09
GMM	1	59	-38474.99	-77303.47	0	0.50
	2	119	-35524.70	-71762.39	0.77	0.06
	3	179	-34872.72	-70817.91	0.51	0.29
	4	239	-34544.54	-70521.04	0.52	0.39
	5	299	-34356.91	-70505.26	0.46	0.45
	6	359	-34252.98	-70656.90	0.44	0.46

6 Discussion

We have introduced novel GHD-GMM and GST-GMM models, which are extensions of the GMMs introduced by Verbeke and Lesaffre (1996) to the generalized hyperbolic and skew-t distributions, respectively, to facilitate heavier tails or asymmetry. Updates are derived for parameter estimation within the EM algorithm framework in the model-based clustering context, which is made feasible by the fact that the generalized hyperbolic distribution can be represented as a normal mean-variance mixture, where the weight follows a GIG distribution. In our extension GMMs, four models were considered (GHD-GMM and GST-GMM under $\beta_{yk} = \mathbf{0}$, GHD-GMM and GST-GMM under $\beta_{\eta k} = \mathbf{0}$) and their performance was compared using simulated and real data. In terms of interpretation, GHD-GMM under $\beta_{\eta k} = \mathbf{0}$ is better than GHD-GMM under $\beta_{yk} = \mathbf{0}$ because the skewness parameters are in the data space and the interpretation of the skewness parameters is clear. However, naturally, models with a minimal number of parameters would be preferable. Hence, in terms of model complexity, GHD-GMM under $\beta_{yk} = \mathbf{0}$ is preferable to GHD-GMM under $\beta_{\eta k} = \mathbf{0}$, because the former model has $K(T - q)$ fewer parameters than the latter.

We believe that this kind of mixture modeling approach for longitudinal data is important in many biostatistical and psychological applications, allowing accurate inference of model parameters and class membership probabilities while adjusting for heterogeneity, heavy tails, and skewness in the data. The proposed GHD-GMM and GST-GMM models have several advantages over Gaussian GMMs. The proposed GHD-GMM, which includes the multivariate skew-t, variance-gamma distribution, multivariate Gaussian distributions, etc., as special or limiting cases, provides flexibility to handle a broader range of multivariate longitudinal data. In the presence of heterogeneity, heavy tails, and skewness in longitudinal data, the proposed method can fit the data considerably better than Gaussian mixtures reducing the risk of extracting latent classes that are merely due to non-normality of the outcomes. When the data are normal, the proposed GHD-GMM can be used to check the reproducibility of a Gaussian GMM solution due to the flexibility of the generalized hyperbolic distribution.

The models proposed herein can be further developed in various ways. First, for the first level of the GMM, only q th order polynomial equations are considered, and so kernel regressions or non-linear regressions could be incorporated into the model. Second, Bayesian mixture modeling may offer researchers an alternative way to handle clustering of longitudinal data due to the popularity and advances in Markov chain Monte Carlo techniques. Finally, it is also worthwhile to consider more general parametric distributions of measurement errors and random effects, such as the coalesced generalized hyperbolic distribution and the multiple scaled generalized hyperbolic distribution (Tortora et al., 2014, 2016).

References

- Aitken, A. C. (1926). A series formula for the roots of algebraic and transcendental equations. *Proceedings of the Royal Society of Edinburgh* 45, 14–22.
- Azzalini, A. and A. Capitanio (1999). Statistical applications of the multivariate skew normal distribution. *Journal of the Royal Statistical Society: Series B* 61(3), 579–602.
- Azzalini, A. and A. D. Valle (1996). The multivariate skew-normal distribution. *Biometrika* 83, 715–726.
- Barndorff-Nielsen, O. and C. Halgreen (1977). Infinite divisibility of the hyperbolic and generalized inverse Gaussian distributions. *Probability Theory and Related Fields* 38(4), 309–311.
- Blæsild, P. (1978). The shape of the generalized inverse Gaussian and hyperbolic distributions. Research Report 37, Department of Theoretical Statistics, Aarhus University, Denmark.
- Böhning, D., E. Dietz, R. Schaub, P. Schlattmann, and B. Lindsay (1994). The distribution of the likelihood ratio for mixtures of densities from the one-parameter exponential family. *Annals of the Institute of Statistical Mathematics* 46(2), 373–388.
- Bouveyron, C. and C. Brunet (2012). Simultaneous model-based clustering and visualization in the Fisher discriminative subspace. *Statistics and Computing* 22(1), 301–324.

- Browne, R. P. and P. D. McNicholas (2015). A mixture of generalized hyperbolic distributions. *Canadian Journal of Statistics* 43(2), 176–198.
- Bryk, A. S. and S. W. Raudenbush (1987). Application of hierarchical linear models to assessing change. *Psychological Bulletin* 101(1), 147.
- Bryk, A. S. and S. W. Raudenbush (1992). *Hierarchical Linear Models: Applications and Data Analysis Methods*. Sage Publications, Inc.
- Dempster, A. P., N. M. Laird, and D. B. Rubin (1977). Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society: Series B* 39(1), 1–38.
- Good, I. J. (1953). The population frequencies of species and the estimation of population parameters. *Biometrika* 40(3-4), 237–264.
- Halgreen, C. (1979). Self-decomposability of the generalized inverse Gaussian and hyperbolic distributions. *Probability Theory and Related Fields* 47(1), 13–17.
- Hubert, L. and P. Arabie (1985). Comparing partitions. *Journal of Classification* 2(1), 193–218.
- Jørgensen, B. (1982). *Statistical Properties of the Generalized Inverse Gaussian Distribution*. New York: Springer-Verlag.
- Kass, R. E. and A. E. Raftery (1995). Bayes factors. *Journal of the American Statistical Association* 90, 773–795.
- Kass, R. E. and L. Wasserman (1995). A reference Bayesian test for nested hypotheses and its relationship to the Schewartz criterion. *Journal of the American Statistical Association* 90(431), 773–795.
- Keribin, C. (2000). Consistent estimation of the order of mixture models. *Sankhyā. The Indian Journal of Statistics. Series A* 62(1), 49–66.
- Laird, N. M. and J. H. Ware (1982). Random-effects models for longitudinal data. *Biometrics* 38(4), 963–974.
- Lee, S. and G. J. McLachlan (2014). Finite mixtures of multivariate skew t-distributions: some recent and new results. *Statistics and Computing* 24(2), 181–202.
- Leroux, B. G. (1992). Mixture models: Theory, geometry and applications. *The Annals of Statistics* 1992, 1350–1360.
- Lin, T.-I., P. H. Wu, G. J. McLachlan, and S. X. Lee (2014). The skew-t factor analysis model. Available from <http://arxiv.org/abs/1310.5336> [accessed on 4 December 2014].
- Lindsay, B. G. (1995). Mixture models: Theory, geometry and applications. In *NSF-CBMS Regional Conference Series in Probability and Statistics*, Volume 5. California: Institute of Mathematical Statistics: Hayward.

- Lu, X. and Y. Huang (2014). Bayesian analysis of nonlinear mixed-effects mixture models for longitudinal data with heterogeneity and skewness. *Statistics in Medicine* 33(16), 2830–2849.
- McArdle, J. J. and D. Epstein (1987). Latent growth curves within developmental structural equation models. *Child Development*, 110–133.
- McLachlan, G. and D. Peel (2000). *Finite Mixture Models*. New York, NY: John Wiley & Sons.
- McNeil, A., R. Frey, and P. Embrechts (2005). *Quantitative Risk Management: Concepts, Techniques and Tools*. Princeton University Press.
- McNicholas, P. D., T. B. Murphy, A. F. McDaid, and D. Frost (2010). Serial and parallel implementations of model-based clustering via parsimonious Gaussian mixture models. *Computational Statistics and Data Analysis* 54(3), 711–723.
- Meng, X.-L. and D. B. Rubin (1993). Maximum likelihood estimation via the ECM algorithm: A general framework. *Biometrika* 80(2), 267–278.
- Meng, X.-L. and D. Van Dyk (1997). The EM algorithm: an old folk-song sung to a fast new tune. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 59(3), 511–567.
- Murray, P. M., R. P. Browne, and P. D. McNicholas (2014). Mixtures of skew-t factor analyzers. *Computational Statistics & Data Analysis* 77, 326 – 335.
- Muthén, B. (2004). Latent variable analysis. *The Sage Handbook of Quantitative Methodology for the Social Sciences*. Thousand Oaks, CA: Sage Publications, 345–68.
- Muthén, B. and T. Asparouhov (2008). Growth mixture modeling: Analysis with non-Gaussian random effects. *Longitudinal Data Analysis*, 143–165.
- Muthén, B. and T. Asparouhov (2015). Growth mixture modeling with non-normal distributions. *Statistics in Medicine* 34(6), 1041–1058.
- Muthén, B. and K. Shedden (1999). Finite mixture modeling with mixture outcomes using the EM algorithm. *Biometrics* 55(2), 463–469.
- Muthén, B. O. and L. K. Muthén (1998–2012). *Mplus User’s Guide*. Seventh Edition. Los Angeles, CA: Muthén and Muthén.
- Nagin, D. (2005). *Group-Based Modeling of Development*. Harvard University Press.
- Nagin, D. S. (1999). Analyzing developmental trajectories: A semiparametric, group-based approach. *Psychological Methods* 4(2), 139.
- Nagin, D. S. and K. C. Land (1993). Age, criminal careers, and population heterogeneity: Specification and estimation of a nonparametric, mixed Poisson model. *Criminology* 31, 327.
- Sahu, S. K., D. K. Dey, and M. D. Branco (2003). A new class of multivariate skew distributions with applications to Bayesian regression models. *Canadian Journal of Statistics* 31(2), 129–150.

- Schwarz, G. (1978). Estimating the dimension of a model. *The Annals of Statistics* 6, 461–464.
- Singer, J. D. and J. B. Willett (2003). *Applied Longitudinal Data Analysis: Modeling Change and Event Occurrence*. Oxford university press.
- Steinley, D. (2004). Properties of the Hubert-Arabie adjusted Rand index. *Psychological Methods* 9, 386–396.
- Tortora, C., B. C. Franczak, R. P. Browne, and P. D. McNicholas (2014). Mixtures of multiple scaled generalized hyperbolic distributions. *arXiv preprint arXiv:1403.2332*.
- Tortora, C., P. D. McNicholas, and R. P. Browne (2016). A mixture of generalized hyperbolic factor analyzers. *Advances in Data Analysis and Classification* 10(4), 423–440.
- Verbeke, G. and E. Lesaffre (1996). A linear mixed-effects model with heterogeneity in the random-effects population. *Journal of the American Statistical Association* 91(433), 217–221.
- Woodbury, M. A. (1950). *Inverting modified matrices*. Statistical Research Group, Memorandum Report 42. Princeton University, Princeton, New Jersey.

A Distribution of $\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, w_{ik}, c_{ik} = 1$

Herein, we give the detailed derivation of the conditional distribution of $\boldsymbol{\eta}_i$ given $\mathbf{y}_i, \mathbf{x}_i, w_{ik}$, and $c_{ik} = 1$, which facilitate the computation of the conditional expectations in the E-step of the EM algorithm. It also serves as a way to estimate the growth factor scores. The joint distribution of $\boldsymbol{\eta}_i$ and $\mathbf{Y}_i$ given $\mathbf{x}_i, w_{ik}$, and $c_{ik} = 1$ is given by

$$\begin{pmatrix} \boldsymbol{\eta}_i \\ \mathbf{Y}_i \end{pmatrix} \Big| \mathbf{x}_i, w_{ik}, c_{ik} = 1 \sim \mathcal{N} \left(\begin{pmatrix} \boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k} \\ \boldsymbol{\Lambda}_y (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k}) \end{pmatrix}, \begin{pmatrix} w_{ik} \boldsymbol{\Psi}_k & w_{ik} \boldsymbol{\Psi}_k \boldsymbol{\Lambda}_y' \\ w_{ik} \boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k & w_{ik} (\boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k \boldsymbol{\Lambda}_y' + \boldsymbol{\Theta}_k) \end{pmatrix} \right).$$

According to the properties of the conditional distribution for multivariate normal variables, the conditional distribution of $\boldsymbol{\eta}_i$ conditional on $\mathbf{y}_i, \mathbf{x}_i, w_{ik}, c_{ik} = 1$ is also a multivariate normal distribution with

$$\begin{aligned} \mathbb{E}(\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, w_{ik}, c_{ik} = 1) &= \boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k} \\ &\quad + \boldsymbol{\Psi}_k \boldsymbol{\Lambda}_y' (\boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k \boldsymbol{\Lambda}_y' + \boldsymbol{\Theta}_k)^{-1} (\mathbf{y}_i - \boldsymbol{\Lambda}_y (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k})), \\ \text{Var}(\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, w_{ik}, c_{ik} = 1) &= w_{ik} \boldsymbol{\Psi}_k - w_{ik} \boldsymbol{\Psi}_k \boldsymbol{\Lambda}_y' (\boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k \boldsymbol{\Lambda}_y' + \boldsymbol{\Theta}_k)^{-1} \boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k. \end{aligned}$$

According to the Woodbury matrix identity (Woodbury, 1950), the covariance matrix for the latent variable $\boldsymbol{\eta}_i$ can be simplified to

$$\text{Var}(\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, w_{ik}, c_{ik} = 1) = w_{ik} (\boldsymbol{\Psi}_k^{-1} + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y)^{-1}.$$

Next, let us simplify the expectation of the latent variable $\boldsymbol{\eta}$:

$$\begin{aligned}
\mathbb{E}(\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, w_{ik}, c_{ik} = 1) \\
&= (\mathbf{I}_T - \boldsymbol{\Psi}_k \boldsymbol{\Lambda}'_y (\boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k \boldsymbol{\Lambda}'_y + \boldsymbol{\Theta}_k)^{-1} \boldsymbol{\Lambda}_y) (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k}) + \boldsymbol{\Psi}_k \boldsymbol{\Lambda}'_y (\boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k \boldsymbol{\Lambda}'_y + \boldsymbol{\Theta}_k)^{-1} \mathbf{y}_i, \\
&= (\boldsymbol{\Psi}_k - \boldsymbol{\Psi}_k \boldsymbol{\Lambda}'_y (\boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k \boldsymbol{\Lambda}'_y + \boldsymbol{\Theta}_k)^{-1} \boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k) \boldsymbol{\Psi}_k^{-1} (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k}) \\
&\quad + \boldsymbol{\Psi}_k \boldsymbol{\Lambda}'_y (\boldsymbol{\Theta}_k^{-1} - \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y (\boldsymbol{\Psi}_k^{-1} + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y)^{-1} \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1}) \mathbf{y}_i, \\
&= (\boldsymbol{\Psi}_k^{-1} + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y)^{-1} \boldsymbol{\Psi}_k^{-1} (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k}) \\
&\quad + (\boldsymbol{\Psi}_k (\boldsymbol{\Psi}_k^{-1} + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y) - \boldsymbol{\Psi}_k \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y) (\boldsymbol{\Psi}_k^{-1} + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y)^{-1} \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i, \\
&= (\boldsymbol{\Psi}_k^{-1} + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y)^{-1} \boldsymbol{\Psi}_k^{-1} (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k}) + (\boldsymbol{\Psi}_k^{-1} + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y)^{-1} \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i, \\
&= (\boldsymbol{\Psi}_k^{-1} + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y)^{-1} (\boldsymbol{\Psi}_k^{-1} (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k}) + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i).
\end{aligned}$$

Finally, we obtain the conditional distribution

$$\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, w_{ik}, c_{ik} = 1 \sim \mathcal{N}(\mathbf{V}_k (\boldsymbol{\Psi}_k^{-1} (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i + w_{ik} \boldsymbol{\beta}_{\eta k}) + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i), w_{ik} \mathbf{V}_k),$$

where $\mathbf{V}_k = (\boldsymbol{\Psi}_k^{-1} + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y)^{-1}$.

B The EM algorithm for Model II

The EM algorithm for Model II was employed for parameter estimation in an analogous fashion to the algorithm for Model I described in Section 4.1. The complete-data comprise the observed outcomes $\mathbf{y}_i$ and covariates $\mathbf{x}_i$, the class membership labels c_{ik} , the latent factors $\boldsymbol{\eta}_i$, and the latent variable w_{ik} , for $i = 1, \dots, n$ and $k = 1, \dots, K$. Therefore, the complete-data log-likelihood is

$$\begin{aligned}
l_c(\vartheta) = \sum_{i=1}^n \sum_{k=1}^K c_{ik} [\log \pi_{ik} + \log \phi(\mathbf{y}_i \mid \boldsymbol{\Lambda}_y \boldsymbol{\eta}_i + w_{ik} \boldsymbol{\beta}_{y k}, w_{ik} \boldsymbol{\Theta}_k) \\
+ \log \phi(\boldsymbol{\eta}_i \mid \boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i, w_{ik} \boldsymbol{\Psi}_k) + \log h(w_{ik} \mid \omega_k, \lambda_k)].
\end{aligned}$$

The E-step requires the computation of the conditional expectations regarding the latent factors $\boldsymbol{\eta}_i$ and the latent variable W_{ik} . Under this formulation,

$$\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, w_{ik}, c_{ik} = 1 \sim \mathcal{N}(\mathbf{V}_k (\boldsymbol{\Psi}_k^{-1} (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}'_y \boldsymbol{\Theta}_k^{-1} (\mathbf{y}_i - w_{ik} \boldsymbol{\beta}_{y k})), w_{ik} \mathbf{V}_k),$$

and the conditional distribution of latent variable W_{ik} given $\mathbf{y}_i, \mathbf{x}_i$, and $c_{ik} = 1$ is given by

$$W_{ik} \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1 \sim \text{GIG}(\psi_k^*, \chi_{ik}, \tilde{\lambda}_k),$$

with $\psi_k^* = \omega_k + \boldsymbol{\beta}'_{y k} \boldsymbol{\Sigma}^{-1}_{y k} \boldsymbol{\beta}_{y k}$, $\chi_{ik} = \omega_k + \delta(\mathbf{y}_i, \boldsymbol{\mu}_k \mid \boldsymbol{\Sigma}_k)$, $\tilde{\lambda}_k = \lambda_k - T/2$, where $\boldsymbol{\mu}_k = \boldsymbol{\Lambda}_y (\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i)$, and $\boldsymbol{\Sigma}_k = \boldsymbol{\Lambda}_y \boldsymbol{\Psi}_k \boldsymbol{\Lambda}'_y + \boldsymbol{\Theta}_k$.

Therefore, we have convenient forms for the following conditional expectations:

$$\begin{aligned}
E_{1ik}^* &:= \mathbb{E}[W_i \mid \mathbf{x}_i, \mathbf{y}_i, c_{ik} = 1] = \sqrt{\frac{\chi_{ik}}{\psi_k^*} \frac{K_{\tilde{\lambda}_k+1}(\sqrt{\psi_k^* \chi_{ik}})}{K_{\tilde{\lambda}_k}(\sqrt{\psi_k^* \chi_{ik}})}}, \\
E_{2ik}^* &:= \mathbb{E}[1/W_i \mid \mathbf{x}_i, \mathbf{y}_i, c_{ik} = 1] = \sqrt{\frac{\psi_k^*}{\chi_{ik}} \frac{K_{\tilde{\lambda}_k+1}(\sqrt{\psi_k^* \chi_{ik}})}{K_{\tilde{\lambda}_k}(\sqrt{\psi_k^* \chi_{ik}})}} - \frac{2\tilde{\lambda}_k}{\chi_{ik}}, \\
E_{3ik}^* &:= \mathbb{E}[\log W_i \mid \mathbf{x}_i, \mathbf{y}_i, c_{ik} = 1] = \log \left(\sqrt{\frac{\chi_{ik}}{\psi_k^*}} \right) + \frac{1}{K_{\tilde{\lambda}_k}(\sqrt{\psi_k^* \chi_{ik}})} \frac{\partial}{\partial \tilde{\lambda}_k} K_{\tilde{\lambda}_k}(\sqrt{\psi_k^* \chi_{ik}}), \\
E_{4ik}^* &:= \mathbb{E}[\boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1] = \mathbf{V}_k(\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1}(\mathbf{y}_i - E_{1ik}^* \boldsymbol{\beta}_{yk})), \\
E_{5ik}^* &:= \mathbb{E}[(1/W_{ik}) \boldsymbol{\eta}_i \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1] = \mathbf{V}_k(E_{2ik}^* (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i) - \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \boldsymbol{\beta}_{yk}), \\
E_{6ik}^* &:= \mathbb{E}[(1/W_{ik}) \boldsymbol{\eta}_i \boldsymbol{\eta}_i' \mid \mathbf{y}_i, \mathbf{x}_i, c_{ik} = 1] = \mathbf{V}_k - \mathbf{V}_k(\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i) \boldsymbol{\beta}_{yk}' \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y \mathbf{V}_k \\
&\quad + E_{2ik}^* \mathbf{V}_k(\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i) (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i)' \mathbf{V}_k, \\
&\quad - \mathbf{V}_k \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \boldsymbol{\beta}_{yk}' (\boldsymbol{\Psi}_k^{-1}(\boldsymbol{\alpha}_k + \boldsymbol{\Gamma}_k \mathbf{x}_i) + \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \mathbf{y}_i)' \mathbf{V}_k + E_{1ik}^* \mathbf{V}_k \boldsymbol{\Lambda}_y' \boldsymbol{\Theta}_k^{-1} \boldsymbol{\beta}_{yk} \boldsymbol{\beta}_{yk}' \boldsymbol{\Theta}_k^{-1} \boldsymbol{\Lambda}_y \mathbf{V}_k.
\end{aligned}$$

At each E-step, the values of E_{1ik}^* to E_{6ik}^* are updated. We also update the value of the class membership variable c_{ik} using

$$p_{ik}^* := \frac{\pi_{ik} f_{\text{GHDT}}(\mathbf{y}_i; \tilde{\lambda}_k, \omega_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k, \boldsymbol{\beta}_{yk})}{\sum_{l=1}^K \pi_{il} f_{\text{GHDT}}(\mathbf{y}_i; \tilde{\lambda}_l, \omega_l, \boldsymbol{\mu}_l, \boldsymbol{\Sigma}_l, \boldsymbol{\beta}_{yl})}.$$

At each M-step, the following model parameters are obtained by maximizing the conditional expected value of $l_c(\vartheta)$ and are updated sequentially. The updates for $\boldsymbol{\pi}_{ik}$, $\boldsymbol{\alpha}_c$, $\boldsymbol{\Gamma}_c$, $\tilde{\lambda}_k$, and ω_k are similar to those used in Section 4.1. We update the skewness parameter $\boldsymbol{\beta}_{yk}$ using

$$\hat{\boldsymbol{\beta}}_{yk} = \frac{\sum_{i=1}^n p_{ik}^* (\mathbf{y}_i - \boldsymbol{\Lambda}_y E_{4ik}^*)}{\sum_{i=1}^n p_{ik}^* E_{1ik}^*},$$

and the measurement error $\boldsymbol{\Theta}_k$ using

$$\begin{aligned}
\hat{\boldsymbol{\Theta}}_k &= \frac{1}{n_k} \sum_{i=1}^n p_{ik}^* \left(E_{2ik}^* \mathbf{y}_i \mathbf{y}_i' - \mathbf{y}_i E_{5ik}^* \boldsymbol{\Lambda}_y' - \mathbf{y}_i \hat{\boldsymbol{\beta}}_{yk}' - \boldsymbol{\Lambda}_y E_{5ik}^* \mathbf{y}_i' + \boldsymbol{\Lambda}_y E_{6ik}^* \boldsymbol{\Lambda}_y' + \boldsymbol{\Lambda}_y E_{4ik}^* \hat{\boldsymbol{\beta}}_{yk}' \right. \\
&\quad \left. - \hat{\boldsymbol{\beta}}_{yk} \mathbf{y}_i' + \hat{\boldsymbol{\beta}}_{yk} E_{4ik}^* \boldsymbol{\Lambda}_y' + E_{1ik}^* \hat{\boldsymbol{\beta}}_{yk} \hat{\boldsymbol{\beta}}_{yk}' \right),
\end{aligned}$$

where $n_k = \sum_{i=1}^n p_{ik}^*$. We update $\mathbf{\Gamma}_k$, $\mathbf{\alpha}_k$, and $\mathbf{\Psi}_k$ sequentially using

$$\begin{aligned}\hat{\mathbf{\Gamma}}_k &= \left\{ \sum_{i=1}^n p_{ik}^* (E_{5ik}^* - E_{2ik}^* \hat{\mathbf{\alpha}}_k) \mathbf{x}_i' \right\} \left\{ \sum_{i=1}^n p_{ik}^* E_{2ik}^* \mathbf{x}_i \mathbf{x}_i' \right\}^{-1}, \\ \hat{\mathbf{\alpha}}_k &= \frac{\sum_{i=1}^n p_{ik}^* (E_{5ik}^* - E_{2ik}^* \hat{\mathbf{\Gamma}}_k \mathbf{x}_i)}{\sum_{i=1}^n p_{ik}^* E_{2ik}^*}, \\ \hat{\mathbf{\Psi}}_k &= \frac{1}{n_k} \sum_{i=1}^n p_{ik}^* \left(E_{6ik}^* - E_{5ik}^* (\hat{\mathbf{\alpha}}_k + \hat{\mathbf{\Gamma}}_k \mathbf{x}_i)' - (\hat{\mathbf{\alpha}}_k + \hat{\mathbf{\Gamma}}_k \mathbf{x}_i) E_{5ik}^{*'} + E_{2ik}^* (\hat{\mathbf{\alpha}}_k + \hat{\mathbf{\Gamma}}_k \mathbf{x}_i) (\hat{\mathbf{\alpha}}_k + \hat{\mathbf{\Gamma}}_k \mathbf{x}_i)' \right).\end{aligned}$$